

Review

# Assessment of Glyphosate Impact on the Agrofood Ecosystem

Yaxin Sang<sup>1</sup>, Juan-Carlos Mejuto<sup>2</sup> , Jianbo Xiao<sup>3,4,\*</sup> and Jesus Simal-Gandara<sup>4,\*</sup> 

<sup>1</sup> College of Food Science and Technology, Hebei Agricultural University, Baoding 071001, China; sangyaxin@sina.com

<sup>2</sup> Department of Physical Chemistry, Faculty of Science, University of Vigo—Ourense Campus, E32004 Ourense, Spain; xmejuto@uvigo.es

<sup>3</sup> State Key Laboratory of Quality Research in Chinese Medicine, Institute of Chinese Medical Sciences, University of Macau, Taipa, Macau, China

<sup>4</sup> Nutrition and Bromatology Group, Department of Analytical and Food Chemistry, Faculty of Food Science and Technology, University of Vigo—Ourense Campus, E32004 Ourense, Spain

\* Correspondence: jianboxiao@uvigo.es (J.X.); jsimal@uvigo.es (J.S.-G.)

**Abstract:** Agro-industries should adopt effective strategies to use agrochemicals such as glyphosate herbicides cautiously in order to protect public health. This entails careful testing and risk assessment of available choices, and also educating farmers and users with mitigation strategies in ecosystem protection and sustainable development. The key to success in this endeavour is using scientific research on biological pest control, organic farming and regulatory control, etc., for new developments in food production and safety, and for environmental protection. Education and research is of paramount importance for food and nutrition security in the shadow of climate change, and their consequences in food production and consumption safety and sustainability. This review, therefore, diagnoses on the use of glyphosate and the associated development of glyphosate-resistant weeds. It also deals with the risk assessment on human health of glyphosate formulations through environment and dietary exposures based on the impact of glyphosate and its metabolite AMPA—(aminomethyl)phosphonic acid—on water and food. All this to setup further conclusions and recommendations on the regulated use of glyphosate and how to mitigate the adverse effects.

**Keywords:** glyphosate use; resistant weeds; AMPA; cancer risks; herbicide research



**Citation:** Sang, Y.; Mejuto, J.-C.; Xiao, J.; Simal-Gandara, J. Assessment of Glyphosate Impact on the Agrofood Ecosystem. *Plants* **2021**, *10*, 405. <https://doi.org/10.3390/plants10020405>

Academic Editor: Fabrizio Araniti

Received: 21 January 2021

Accepted: 17 February 2021

Published: 20 February 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.




**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Glyphosate (*N*-phosphonomethylglycine; Figure 1a) is an aminophosphonate. This compound is typically used as a broad-spectrum herbicide and is absorbed by plant leaves. Glyphosate, discovered in the 1970s, was registered in more than 130 countries [1], and the use of glyphosate-based herbicides increased 100 times since then [2]. Genetically engineered herbicide-tolerant (GEHT) crops have considerably facilitated weed management in cotton, soybean, and maize [3–5]. However, they have also caused the emergence of glyphosate-resistant weed phenotypes [3–8]. The incorporation of additional herbicides into spraying programs [6,7] has caused herbicide per hectare on crops with GEHT varieties to escalate in this century [5,8,9]. This upward trend is expected to result in heavier environmental loads and increased human exposure to herbicides, including glyphosate and its main metabolite, aminomethylphosphonic acid (AMPA; Figure 1b), and to the adjuvants contained in its formulations. Weed management should face resistance before it happens [10]. It is key to promote changing crops or crop rotations against herbicide resistant (HR) weeds with effective herbicides [11]. Non-herbicidal alternatives (natural products, selective herbicides, mechanical controls, etc.) need to be added to satisfy the reduced efficacy of herbicides [12].

Figure 1c shows the degradation pathway for glyphosate in soil [13]. Although leaching of glyphosate is very unlikely due to its high soil sorption, depending on the type of soil,

it can move to ground and surface waters through leaching and runoff [14]. Human exposure to urban sources of glyphosate should be considered too. Although some nonselective (broad spectrum) herbicides for both urban and home use in emerging countries contain glyphosate at low levels—and pose little risk of acute toxic exposure as a result [15]—those used in developing countries contain higher levels of this compound, or are mixed without official control. Dietary exposure in areas lacking residue information can be assessed from data for areas where glyphosate use and residues have been accurately determined [16].



**Figure 1.** Structural formula of glyphosate (a) and AMPA –(aminomethyl)phosphonic acid– (b), together with degradation pathway for glyphosate in soil (c).

The glyphosate mechanism as herbicidal involves inhibition of 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS), which interferes with phenylalanine, tyrosine, and tryptophan synthesis. Unlike plants and some microorganisms, mammals have no EPSPS, which is in principle an advantage safety-wise [17]. However, glyphosate herbicides are highly controversial in toxicological and environmental terms. This review, therefore, diagnoses on the use of glyphosate and the associated development of glyphosate-resistant weeds. It also deals with the risk assessment on human health of glyphosate formulations through environment and dietary exposures based on the impact of glyphosate and its

metabolite AMPA on water and food. All this to setup further conclusions and recommendations on the regulated use of glyphosate and how to mitigate the adverse effects in the below selected sections. The literature search was done following the guidelines included in the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guide (PRISMA). Accordingly, a search was carried out on the following databases: Cochrane library, Embase, Medline, Cinahl, Scopus, Sci-Finder, and Web of Science. Scientific reports included in the study were obtained using the following search terms: “(glyphosate) AND (keywords of the different sections below)”. Common inclusion/exclusion criteria for references selection were based on publication date, mainly in the past 20 years, of both scientific-based papers and technical reports, with study designs, interests, and world overall coverage, applicable to all research questions below.

## 2. Glyphosate-Resistant Weeds

The international database on herbicide resistance [18] contains more than 510 studies. The best resistance mainframe is based on prevention and on detection with regular appraisal of herbicides-treated fields [19]. There are various methods to detect resistance with tests in the field and with bioassays in greenhouses and laboratories [20]; for example, hybridization between *A. palmeri* and *A. spinosus* occurred with frequencies in the field studies ranging from <0.01% to 0.4%, and 1.4% in greenhouse crosses [21]. Non-target-site resistance (NTSR) to herbicides in weeds can be conferred as a result of the alteration of one or more physiological processes, including herbicide absorption, translocation, sequestration, and metabolism. The mechanisms of NTSR are generally more complex to decipher than target-site resistance (TSR) and can impart cross-resistance to herbicides with different modes of action. Metabolism-based NTSR has been reported in many agriculturally important weeds, although reduced translocation and sequestration of herbicides has also been found in some weeds [22,23]. Crossed resistance is when the plant developed resistance to an herbicide, which permits to resist herbicides with the same action mode [24]. Multiple resistance is when a plant has one or several mechanisms of resistance to herbicides with distinct action modes. The selection pressure of an herbicide is then capable to select resistant plant biotypes depending on the herbicide treatment type, its formulation, application frequency, and the biological characteristics of the weed and the crop [25–28]. Examples of glyphosate-resistant weeds and their locations can be found in Table 1.

**Table 1.** Example of glyphosate-resistant weeds and their locations, extended from reference [29].

Weed	Location
<i>Amaranthus palmeri</i>	United States
<i>Amaranthus tuberculatus</i>	United States
<i>Ambrosia artemisiifolia</i>	United States
<i>Ambrosia trifida</i>	United States
<i>Conyza bonariensis</i>	United States, Brazil, Argentina
<i>Conyza canadensis</i>	United States
<i>Euphorbia heterophylla</i>	Brazil
<i>Lolium perenne</i>	United States, Brazil, Australia
<i>Sorghum halepense</i>	United States, Argentina

The problem is compounded by non-target site multiple resistances in grasses, as is the case of *Lolium rigidum* and *Alopecurus myosuroides* [12,26]. In addition, the growing expansion of multiple resistance to broadleaf weeds is bound to worsen things in the future. Managing non-target site resistance is difficult owing to the many unpredictable resistance patterns against which rotating herbicide sites of action may be ineffective. Herbicides are

the main means for weed control in developed countries, but they should be used more sustainably [30]. This entails not only using improved herbicide mixtures and rotations, but still adopting intensive integrated weed management programs including effective mechanical and cultural strategies [31]. A pressing need therefore exists for economic incentives to the search for new, safer, and more effective herbicides [4].

Developing herbicide-resistance crop traits may grant the use of old herbicides in new ways through tailored mixtures efficiently avoiding multiple resistance [32,33]. Indeed, the use of different genetic engineering techniques as RNA interference (RNAi) [3,32,34–38], chimeric RNA/DNA oligonucleotides [39], and gene-editing techniques (GM), such as CRISPR/Cas9 or CRISPR/Cpf1 [40–48] technology, might be useful for this purpose. The best approach to prevent resistant weeds is to use a combined weed management, and herbicides will likely be partly replaced with new technologies such as, among others, research on crop allelopathy [49–56] and engineering of microbial control agents [57–61]. Progress in these technologies is expected to allow methods for weed control to be used in an integrated manner with the aim of maximizing diversity in weed control and minimizing resistance. Applying evolutionary principles to agricultural settings is essential to properly understand the system-wide effects of herbicide selection intensity [62]. Although the main driver of herbicide resistance is the selection pressure of management, further knowledge of the scientific bases of herbicide resistance at the genetic and cellular levels needs to be developed [63–67]. The causes and dynamics of resistance expansion might be elucidated by assessing the flexibility of certain alleles involved in herbicide resistance [68]. The capability of resistant weeds to prevail, replicate, and selectively infest habitats depends on the degree of vigour of the particular resistant gene [69]. The effects of the environment on resistant plants in cropping conditions could thus reduce the heritability and frequencies of resistance alleles with time [70].

Research in this field should also address the effects of climate change on the expansion of herbicide resistance. According to Renton [71], spatially computational models could be of help in this context by providing powerful indicators on how genetics, plant biology, population structure, environmental conditions, and management strategies connect to shape weed resistance dynamics. The use of scarcely diverse practices drove to a fast increment in multiple-resistance weeds with upgraded abilities for herbicide metabolism worldwide [72]. Elucidating herbicide metabolic pathways could help re-classify and re-rank the risks of herbicide resistance and hence enable the adoption of more effective herbicide rotations, such as those based on both site of action and metabolic pathway. Processes associated with climatic changes, such as elevated temperatures, can strongly affect weed control efficiency. For example, responses of several grass weed populations to herbicides that inhibit acetyl-CoA carboxylase (ACCCase) were examined under different temperature regimes [73].

### *Mitigation Strategies*

Weed resistance management has three planks: rotate modes of action to reduce selection pressure, incorporate non-chemical practices and control weed seed set, together with asexual reproduction (rhizomes, stolons, etc.). Effective weed control needs to discern weeds biology, prevents with weed seed production, plant into weed-free fields, grow weed-free seed, and inspect lands regularly. There is also a need to adopt numerous herbicide action mechanisms active versus damaging weeds, spread herbicide estimate at selected weed extents, and highlight growing conditions that put an end to weeds by crop competition. Further, it is useful to practice mechanical and biological executive strategies, avoid field-to-field and within-field migration of weed vegetative propagules, regulate weed seed to avoid a reinforcement of the weed seed-stock, and preclude an invasion of weeds into land by controlling ground boundaries. All these diverse approaches to managing herbicide resistance need to be incorporated into weed management [74–78].

This will be beneficial in managing resistance in the long term, together with mathematical simulations proving that mixtures magnify herbicide efficacy, choice array of soil-applied herbicides, and postpone herbicide resistance growth in weeds. It shows that extension efforts rotating herbicide mixtures give vision to guide the progression of weed resistance [69]. Multiple modes of action (MOAs) for weed control are important for managing herbicide resistance and enabling no-till farming practices that help to sequester greenhouse gases, but discovering new herbicide MOAs has been a challenge for the industry [76].

According to the International Assessment of Agricultural Knowledge, Science and Technology for Development [79] (IAASTD, 2008), agricultural development has focused on increasing farm-level yield, more than on consolidating effects on biodiversity and the liaison of agriculture with climate change. Increased attention needs to be directed to build up soil fertility and to sustain agricultural production, with a focus also on protection of biodiversity. Agro-ecology refers to treating agricultural ecosystems as ecosystems, and can enable a successful transition to more sustainable farming and food systems [80].

Moreover, in recent decades, studies were performed looking for alternatives to glyphosate. There is a rise in efficacy tests using different natural (or even modified) allelo-chemicals obtained from essential oils for pest-control [81–96], as well as in their use as herbicides [97–112].

### 3. Impact of Glyphosate and Its Metabolite AMPA on Water Streams

Glyphosate residues raised perception of its adverse effects on human health, soil, and aquatic ecosystems [113]. Some microorganisms in soil and water can degrade this compound [114]. The major metabolite of glyphosate is aminomethylphosphonic acid (AMPA; Table 2).

**Table 2.** Properties of glyphosate and its metabolite AMPA.

Common Name	Glyphosate	AMPA
Chemical name	<i>N</i> -phosphonomethylglycine	Aminomethylphosphonic acid
CAS number	1071-83-6	1066-51-9
Molecular formula	C <sub>3</sub> H <sub>8</sub> NO <sub>5</sub> P	CH <sub>6</sub> NO <sub>3</sub> P
Exact mass	169.01 g mol <sup>-1</sup>	111.01 g mol <sup>-1</sup>
Vapour pressure (25 °C)	1.31 × 10 <sup>-5</sup> Pa	8.44 × 10 <sup>-4</sup> Pa
Henry's law volatility constant (25 °C)	2.1 × 10 <sup>-7</sup> Pa m <sup>3</sup> mol <sup>-1</sup>	2.6 × 10 <sup>-3</sup> Pa m <sup>3</sup> mol <sup>-1</sup>
Solubility in water (20 °C)	10.5 g L <sup>-1</sup>	1467 g L <sup>-1</sup>
Partition coefficient (log P <sub>ow</sub> ) (20 °C)	-3.2	-1.6
Solid/water distribution coefficient (K <sub>d</sub> )	5.3–900 L kg <sup>-1</sup>	15–1554 L kg <sup>-1</sup>
Soil organic carbon normalized adsorption coefficient (K <sub>oc</sub> )	884–60,000 L kg <sup>-1</sup>	1160–24,800 L kg <sup>-1</sup>
Half-life (DT <sub>50</sub> ) in soil	1–197 days	23–958 days
DT <sub>90</sub> in soil	40–280 days	Unknown

Glyphosate can flow throughout soil, and reach surface and ground waters [14,115–117]. Although sorption and degradation are affected by many factors that might be expected to affect glyphosate mobility in soils, glyphosate leaching seems mainly determined by soil structure and rainfall. Limited leaching has been observed in non-structured sandy soils, while subsurface leaching to drainage systems was observed in a structured soil with preferential flow in macropores, but only when high rainfall followed glyphosate application [14]. The time needed for glyphosate in river water to be eliminated by 50% (i.e., DT<sub>50</sub>) has been found to range from 13.8 to 301 days, which is suggestive of moderate to high persistence [118]. Rivers are influential environments with a fundamental action

in xenobiotic mitigation [119]. Thus, river water usually transports nutrients, organic matter, and pollutants that can severely constrain microbial growth [120]. Biofilms, which are network structures with a wide range of microbes, contribute to transform xenobiotics through co-metabolism and mineralization. Glyphosate can be utilized to obtain phosphorus by microorganisms such as bacteria and fungi in biofilms [121,122]. The compound is cleaved through the carbon–phosphorus lyase (C–P lyase) route, which comprises consecutive enzyme-catalysed reactions including phosphonate activation and C–P bond break [123]. The genes enciphering the enzymes are amassed into the phosphonate operon, which occurs widely amongst bacteria. However, the effects of phosphorus on glyphosate degradation have been investigated in segregated microbial strains [124] rather than in natural biofilms [125].

AMPA is a metabolite from glyphosate and from aminopolyphosphonate, which is applied in detergents, flame-retardants, and anticorrosive products [126]. As found by meta-analysis [127], glyphosate and AMPA are concomitantly present in water, with levels evolving in the same way at positively correlated concentrations, since glyphosate can be rapidly converted to AMPA. Battaglin et al. [128] detected glyphosate, but no AMPA in 2.3% of 3732 water and sediment specimens. Moreover, Struger et al. [129] found the parent compound and its metabolite to co-occur at probability  $p$  value = 0.76 in Canadian rivers. The concomitance of glyphosate and AMPA in groundwater suggests partial mineralization of the former under the influence of anthropic activities or environmental conditions. Glyphosate is converted into AMPA largely in eutrophic water than it is in P-poor water [127]. Glyphosate and AMPA have been identified at oligotrophic sites and found to be completely absent from upstream communities under low glyphosate–low phosphorus conditions. Glyphosate can have from low to high persistence in soils with aerobic conditions ( $DT_{50}$  2.8–500.3 days), and high persistence in anaerobic soils ( $DT_{50}$  135–1000 days) [118]. Glyphosate persistence in water sediments is moderate to high ( $DT_{50}$  13.82–301 days) [118]. Phosphorous from glyphosate and AMPA is very low accounting for <0.17% of total P. The degree of contamination of surface water with glyphosate depends largely on two factors, namely: (a) herbicide level, affected by soil biodegradation and sorption; and (b) phosphorus availability in the water, reducing its degradation. This should be considered in the evaluation of the environmental risks of glyphosate and AMPA present in surface waters [121,127,130–137].

As some people might use surface-water for drinking and preparation of food, it was assumed that it was consumed untreated. Based on the median (0.03 µg/L) and 99th centile (302 µg/L) concentrations found in surface-water, oral doses were 0.00000086 and 0.0043 mg/kg b.m./d. These exposures are considerably less than the acceptable daily intakes (ADIs) and present de minimis risk [138]. Glyphosate can be bound to divalent and trivalent cations in the soil and water, but there is risk to aquatic life joined to the residues of free available glyphosate in water [139]. The hazard quotients (HQs) obtained in sediments and the repercussion over benthic creatures were of 1.4–6.7, proposing risk for sediment dwellers. Annett et al. [140] and Thompson et al. [141] estimate HQs higher than one for fish and for aquatic microorganisms, together with invertebrates and amphibians. The HQs for AMPA advice it is not threatening. Glyphosate seems to be critical where intensive agriculture is practiced, since it involves increased use of fertilizers, plant growth regulators, and pesticides and mechanised agriculture [130,139,140,142].

#### *Mitigation Strategies*

Glyphosate residues are linked to its applications calendar and soil inputs; thus, the need to reduce overspray, but also to focus on its relative mobility/persistence [143]. Residual levels in sediments and water streams may be diluted with the upper part of the basin devoted to woodland, because it is not used in woodlands, and woodlands themselves help to reduce and prevent diffuse pollution [144]. There is a need to develop sediment quality guidelines for such contaminants [145]. The actions to restrict them in surface and groundwater below current water quality standards can be at the exploitation level

(collection and treatment of wash water for sprayers or engines . . . ), at field plot level (reduction of the dose, application date according to the weather . . . ), and at catchment level (vegetated buffer strips, orientation of the crop rows . . . ). Artificial wetlands have to be implemented in addition to local action, such as a pesticide reduction plan [146]. The most commonly used mitigation techniques to prevent pesticide input into water bodies include edge-of-field and riparian buffer strips, vegetated ditches, and constructed wetlands. It has been identified that removal of pesticides is highly variable, and generally increases with increasing value of  $K_{OC}$ , but the relationship is not strong [147]. All these undertakings should help with the mandatory regulations for Sustainable Use of Pesticides Directive, which include each EU member state having a National Action Plan on the reduction in the use of pesticides, buffer zones, prevention of contamination of watercourses, etc.

#### 4. Glyphosate-Based Herbicides and Cancer Risks

The mechanisms of action of glyphosate herbicides involve endocrine or microbiome disruption (Table 3).

**Table 3.** Some recent studies of glyphosate-based formulations' toxicity involving various indicators such as organisms that can be affected and the main results at tested concentrations.

Herbicide Formulation	Test-Organism	Endpoint	Results	Tested Concentrations	Ref
Glyphosate	<i>Geotrichum candidum</i> , <i>Lactococcus lactis</i> subsp. <i>Cremoris</i> ; <i>Lactobacillus delbrueckii</i> subsp. <i>bulgaricus</i>	Microbial growth assay	Inhibition of microbial growth by the commercial product Roundup; microbiocidal effect at concentrations below those recommended for agricultural use of the commercial product Roundup; no significant toxicity of the active ingredient (glyphosate) on any of the microorganisms	0.1, 1, 10, 100, 1000, 10,000 ppm	[148]
Glyphosate	Tadpoles of wood frog ( <i>Rana sylvatica</i> or <i>Sylvaticus lithobates</i> ), leopard frog ( <i>Rana pipiens</i> or L.) and American toad ( <i>Bufo americanus</i> or <i>Anaxyrus americanus</i> )	Acute toxicity assay	Significant induction of morphological alterations in tadpoles of the three species; exposure to glyphosate altered tadpole tail size in wood and leopard frogs at all tested concentrations	0, 1, 2, or 3 mg acid equivalents [a.e.]/L of Roundup Original MAX	[149]
Glyphosate	Roots from the smooth hawksbeard ( <i>Crepis capillaris</i> L.); polychromatic erythrocytes of the bone marrow of C57BL rat	Chromosome aberration assay; micronucleus assay	No induction of genotoxic and/or mutagenic effects on any of the species	<i>Crepis capillaris</i> : 0.05, 0.1, 0.5, 1%; erythrocytes: doses inferior to half the LD <sub>50</sub> (1080 mg/Kg)	[150]
Glyphosate	Female Wistar rats	Acute toxicity assay; teratogenicity assay	High mortality index of females treated with the highest concentration of the commercial product Roundup; increased dose-response of foetal skeletal alterations	500, 750, 1000 mg kg <sup>-1</sup>	[151]

Table 3. Cont.

Herbicide Formulation	Test-Organism	Endpoint	Results	Tested Concentrations	Ref
Glyphosate	Human lymphocytes	Comet assay; FISH; lipid peroxidation assay–TBARS	Significantly increased DNA migration at 580 $\mu\text{g mL}^{-1}$ ; significantly increased comet tail intensity at 92.8 $\mu\text{g mL}^{-1}$ ; increased DNA damage in the presence of S9; increased frequency of micronuclei, nuclear buds and nucleoplasmic bridges, without S9; significantly increased nuclear instability at the highest concentration tested with S9; significantly increased dose–response of TBARS levels	0.5, 2.91, 3.5, 92.8, 580 $\mu\text{g mL}^{-1}$	[152]
Glyphosate; 2,4-D	Algae and 25 species of aquatic animals	Acute toxicity assay	No reduction in periphyton biomass by either herbicide; no strong impact of 2,4-D on the aquatic community; strong impact of glyphosate on the aquatic community (significantly decreased species richness)	0, 1, 2, or 3 mg acid equivalents [a.e.]/L of Roundup® Original MAX	[149]
Glyphosate/As As/Cu	Soil nematode <i>Caenorhabditis elegans</i>	Heat Shock Protein Response, Reproduction and Locomotory behaviour (head thrashing)	Responses in locomotory behaviour (head thrashing), reproduction, and heat shock protein expression had been observed.	Sublethal 24-h exposures of 1/1000, 1/100 and 1/10 of the LC50	[153]
Glyphosate; Terbutylazine	Human lymphocytes	Cytome FISH	Glyphosate concentrations above 3.5 $\mu\text{g mL}^{-1}$ increased the frequencies of micronuclei, nuclear buds and nucleoplasmic bridges in treated cells without inducing centromeric signals; terbutylazine at concentrations above 0.008 $\mu\text{g mL}^{-1}$ increased the frequency of micronuclei hybridized with centromeric probe and of nuclear buds with centromeric signals in the presence of S9	0.5, 2.91, 3.50, 92.8, 580 $\mu\text{g mL}^{-1}$ (glyphosate); 0.00058, 0.0008, 0.008, 25, 156.5 $\mu\text{g mL}^{-1}$ (terbutylazine)	[154]
Glyphosate	Earthworms <i>Pontoscolex corethrurus</i> <i>Amyntas corticis</i>	Toxicity assay	Coffee plantations with regular applications of Glyphosate over the preceding 22 years. Control plantations had received no herbicides over the preceding 7 years. The earthworm species found in plots with no treatment were <i>Pontoscolex corethrurus</i> (99%) and <i>Amyntas corticis</i> (1%), while <i>A. corticis</i> was absent in plots that had been treated.	Manufacturer's recommendations	[155]



Table 3. Cont.

Herbicide Formulation	Test-Organism	Endpoint	Results	Tested Concentrations	Ref
Glyphosate	<i>Allium cepa</i>	Cytotoxic evaluation Cytogenotoxic effects	Exposure to glyphosate of <i>A. cepa</i> meristematic cells induces diverse types of chromosomal anomalies in demonstrates that it has a highly cytogenotoxic effect for any of the concentrations used.	5, 10, 15, 25, 30 mg L <sup>-1</sup>	[156]
Glyphosate, alkyphenolpolyglycol ether	Neotropical fish <i>Piaractus mesopotamicus</i> , <i>Phallocerus caudimaculatus</i> , <i>Hyphessobrycon eques</i> , <i>Brachydanio rerio</i>	Toxicity assay, Histopathological effects	The histopathological effects caused by glyphosate exposure on gills, liver, and kidneys are reversible, except for the liver necrosis on <i>P. caudimaculatus</i> . <i>H. eques</i> , <i>P. caudimaculatus</i> , and <i>P. mesopotamicus</i> present great potential to be used as standard organisms for herbicides monitoring and the use of glyphosate without surfactant addition is enough to cause histological alterations on <i>H. eques</i> and <i>P. caudimaculatus</i>	Manufacturer's recommendations Rodeo® Rodeo®+0.5%Aterbane®BR Rodeo®+1.0%Aterbane®BR	[157]
Glyphosate/P	<i>Capsicum annuum</i> Inoculated and non-inoculated with <i>Glomus mosseae</i> or <i>Glomus intraradices</i>	Accumulation of shikimic acid in mycorrhized <i>Capsicum annuum</i> L.	Remobilization of glyphosate residues in the soil by the addition of phosphate should be considered a serious problem for crops in treated soils. The mycorrhization increases the pepper plant's tolerance to high glyphosate concentration in the substrate, and may allow support to this stress condition	Manufacturer's recommendations RoundUp®	[158]
Glyphosate	<i>Misgurnus anguillicaudatus</i>	Toxic assay	Glyphosate represent a potential risk to loach through inhibiting proliferation of diploid and triploid cell lines and induces micronuclei and apoptosis.	80, 240, 400, 560, 720, 880, 1040 mg/L	[159]
Glyphosate	<i>Pouteria torta</i>	Changes in the biological performance	In response to glyphosate, <i>P. torta</i> exhibited reductions in photosynthesis and chloroplastid pigment content, as well as accumulation of shikimic acid and the occurrence of chlorosis and necrosis. These changes demonstrate use as a bioindicator of this herbicide.	25, 50, 100, 200, 400, 800, 1200 g a.e. ha <sup>-1</sup>	[160]

Table 3. Cont.

Herbicide Formulation	Test-Organism	Endpoint	Results	Tested Concentrations	Ref
Glyphosate	<i>Crassostrea gigas</i>	Embryo-larval development and metamorphosis	Embryo-larval development of <i>C. gigas</i> was more sensitive to glyphosate-based herbicides compared to various endpoints studied in regulatory model organisms, and embryos and D-shaped larvae were more sensitive compared to pediveliger larvae.	0.1 to 100,000 µg L <sup>-1</sup> RoundUp®	[161]
Atrazine, Glyphosate	<i>Biophalaria glabrata</i>	Cytotoxic assay	Results indicated that those atrazine and glyphosate herbicides may be considered to be highly genotoxic agents	-	[162]

International regulatory agencies typically classify substances according to their dose–response relationships, thereby overlooking non-monotonic carcinogenic issues in glyphosate. Usually, toxicological data were obtained with an incomplete judgement of the outcome of hormone imitation and the microbiome. Some agencies, including US EPA—United States Environmental Protection Agency, EFSA—European Food Safety Authority, and Canada’s Pest Management Regulatory Agency, are reviewing studies on glyphosate’s effects on human health and species at risk to protect farm workers, food safety, and endangered species. The present decision for the parts of the assessment that are complete, which stands until the next review, shows that can still be recommended when used according to instructions on the label. In March 2015, IARC -International Agency for Research on Cancer—categorized glyphosate as “probably carcinogenic to humans” (Group 2A) based mainly on research demonstrating that ‘there was limited evidence of carcinogenicity in humans’, mostly from agricultural workers, but also concluded that there was ‘sufficient evidence of carcinogenicity in experimental animals’ [163–167]. IARC also concluded that there was evidences for genotoxicity, both for active ingredient and formulations. According to De Roos et al. [168] and further [169], the main conclusion is that “The most reliable approach will be to reanalyse the data after more cases accumulate, both to assess whether the association with myeloma persists and to further evaluate confounding and selection bias using a larger case group to support analyses”. With the same prospective cohort study, Andreotti et al. [170] concluded that there was some evidence of increased risk of acute myeloid leukaemia (AML) among the highest exposed group that requires confirmation. Their effects caused by disturbance of cell–cycle management might also be important for cancer and non-cancer health outcomes [171,172].

There has been a strong controversy over the use of this herbicide and the detection of potential toxic consequences of pure glyphosate itself and glyphosate-based herbicide ingredients that might have a synergistic effect, such as the surfactants used. The surfactants in glyphosate formulations (especially polyoxyethylene, POE-15) are major contributors to DNA damage caused by glyphosate-based herbicides (GBH). Such surfactants have demonstrated to alter mitochondrial function [173] and are also deleterious to human embryonic and placental cells at concentrations around 1 ppm [174]. Richard et al. [175] found Roundup formulation with surfactants to be more than double active than glyphosate alone at producing lethal danger in human placental cells. Moreover, Guilherme et al. [176] found increased numbers of double-strand breaks (DSB) with the comet assay and micronucleus (MN) lesions in eels after exposure to environmental levels of Roundup (0.05 ppm). Therefore, there is a need to regulate the use of GBH in mixtures. In this regards, it was found that glyphosate and its overall formula show genotoxicity in vivo and in vitro [177]. Polyethoxylated tallow amine (POEA) and other surfactants were also

found to be toxic [178]. EU prohibited POEAs in glyphosate formulas, but EPA permitted it at up to 25% (w/w). IARC [179] evidenced that glyphosate formulas produce non-Hodgkin lymphoma (NHL), according to epidemiological research. EPA focused on the Agricultural Health Study [170], finding no relationship of glyphosate with NHL incidence in U.S. applicators. Leon et al. [180] obtained a meta-hazard ratio of 1.36 between diffuse large B-cell lymphoma and glyphosate. Zhang et al. [181] detected that most exposed users had a 41% higher risk of NHL. EPA's Office of Research and Development (ORD) concluded that glyphosate is "likely to be carcinogenic" or "suggestive of carcinogenicity" [182,183].

The results depend on the particular cell, application, and design specifications. Thus, the overall formula exhibits a linear dose-dependent response indicating that toxicity from the adjuvants is monotonic [184]. Gasnier et al. [185] found glyphosate at concentrations below 0.05% to have a non-linear effect on oestrogen receptor-reporter transfected HepG2 cells and the full formulation to linearly reduce androgen receptor-induced transcription with low concentrations. Testosterone-producing Leydig cells afford an alternative model for endocrine disturbance in vivo and ex vivo [148]. Walsh et al. [186] detected disturbed progesterone yield, but only with the full formula, which altered puberty progression and reduced serum testosterone in pre-pubertal Wistar rats at 5 mg kg<sup>-1</sup> day<sup>-1</sup>. Some authors [187–189] recognized the non-monotonicity of glyphosate itself on a human hormone-reliant cell line of breast cancer. They found the effect to be propitiated by the oestrogen feedback and hindered by inclusion of an oestrogen receptor antagonist. Armiliato et al. [190] reported elevated expression of steroidogenic factor-1 and oocyte rise in zebra fish in the microgram-per-litre range in water. However, no significant association with endocrine disruption was found in trout. Glyphosate did not elevate vitellogenin plasma concentrations in young rainbow trout [191]. Gandhi et al. [192] found environmental concentrations of glyphosate in water to alter behaviours such as movement frequency in larval amphibians. If this was the result of a non-monotonic mechanism, then, even very low doses may have some effect on the nervous system. Thus, low levels of glyphosate—even those below regulatory limits—may boost human carcinogenesis through endocrine mimicry.

AMPA is also seemingly genotoxic. Guilherme et al. [193] found 11.8 µg L<sup>-1</sup> concentrations of the glyphosate metabolite to induce DSB in an eel model. In addition, Mañas et al. [194] found AMPA to induce cracks at 2.5 mM in human lymphocyte cultures and in mice. Calculations of total exposure to this degradation product should therefore include residues potentially present in organisms and the environment. Some animal cancer works were proving increased risks of hemangiosarcoma, renal tubule carcinoma, and pancreatic cell islet adenoma, together with skin tumour build up in a mouse model [165]. If both the parent molecule and its metabolite are carcinogenic, then the risk cannot be accurately assessed with the standard Paracelsian dose-response model.

The exposure scenario assessment reflects short-term incidental oral exposure to glyphosate-treated park areas (post-application exposure). The short-term assessment is protective of intermediate-term exposure, and the life-stages selected for aggregate risk assessment are considered protective for the exposures and risks for any other potentially exposed life-stage, since the resulting margins of exposure (MOE), which are Incidental oral NOAEL (No observed adverse effect level)/Residential post-application total exposure, are of 2,200,000 for adults, and of 640 for children 1 to <2 years old [163]. For a chemical substance with health thresholds (i.e., not genotoxic and not carcinogenic), a MOE ≥ 100 is generally considered to be protective. Instead, for genotoxic and carcinogenic compounds, in general a MOE ≥ 10,000 is considered to be protective.

### *Mitigation Strategies*

Glyphosate's use is four times higher than atrazine (the second pesticide in the list) [195]. For this reason, the main mitigation strategy is the control of its use. There is typically a time lag of decades between exposure to a carcinogen and elevated cancer rates, and glyphosate use has skyrocketed over the past 10–15 years, the full effects of glyphosate's rising use remain to be discovered [196].

## **5. Risk Assessment of Glyphosate through Environment and Dietary Exposures**

Pesticides, such as glyphosate, are evaluated periodically for changes. In the case of glyphosate, there is a strong controversy with the results obtained by different authors and the conclusions of different agencies. These contradictions have led glyphosate use to being banned or strictly regulated in some countries. Based on dietary risk assessment [197], glyphosate's NOAEL is 100 mg/kg·day. It is 1114 times higher than the exposure for the US population, and 438 times higher than that of kids between 1 and 2 years [163]. Acquavella et al. [198–200] found a maximum systemic dose for farmers of 0.004 mg/kg. McGuire et al. [201] monitored glyphosate in the urine of breastfeeding women. It was found that only 20% of dietary glyphosate is available, and most of it is excreted in the urine [138,202]. Stephenson and Harris [203,204], considering food processing on glyphosate residues, reduced estimated exposures by 67-fold. Drinking water exposures can be estimated with models based on physical properties of the pesticide, its use, and environmental variables as soil type and rain [205]. Glyphosate is detected in the urine of farm and non-farm family members, kids included, with analogue exposure [206,207]. Glyphosate was also found in human blood [208]. EPA's safe maximum of glyphosate exposure is six times that of Europe's [209]. EPA's estimation of children exposure to glyphosate is higher than the maximum level suggested [210,211]. All of the environmental exposure studies at Connolly et al. [212] had mean/median levels that were 2% of the ADI or less, and the maximum concentrations found in these studies were all less than 6%. One study on residential exposures showed median and maximum values that were 49% and 53% of the ADI, respectively, while another study reported a maximum value was 87% of EFSA's ADI; these studies with the highest percentage compared to the ADI are studies that were outside Europe and involved aerial spraying.

### *Mitigation Strategies*

The key is post-approval monitoring [213]. Such control will track tendencies, identify inflection points, and measure the efficacy of past risk-mitigation assays. Many of the inert ingredients in formulated pesticides are themselves toxic [197,214,215], or help the active ingredient to endure in biological systems. It could be simple to estimate a 5-year rolling average number of herbicides kill units to bear a crop to harvest [216]. If the kill unit begins to slope upward, a new pest management action is necessary. Risks could be reduced by commanding identical pre-harvest intervals on herbicides leading to higher residues, similar number of application rates, compulsory resistance management, and decreased tolerances to prohibit applications. Plans are necessary for these strategies to become real [211,217,218].

## **6. Challenges and Opportunities for Herbicide Research and Development**

The expected growth in the global population will inevitably have to be met by increasing food production. Although the arable land area seemingly remains stable [219], the increasing loss by effect of urbanization and climate change must be considered. Historically, cities have grown in places where good arable land was available [220]. This forced production sectors to increase their output through, for example, efficient weed control. The decreasing variety of effective herbicides and modes of action has had an adverse impact on plant diversity and is hampering sustainable weed management [221]. The introduction of new weed species through international transport of goods is posing additional problems. In addition, the increasingly frequent occurrence of environmental

extremes [222] may affect weeds disparately and alter crop responses. As shown by the giant hogweed (*Heracleum mantegazzianum* Sommier & Levier), invasive weeds can pose serious problems to European authorities [223].

Screenings for structure–activity relationships and virtual screening technologies [97] are helpful to select compounds with diverse structures but identical performance. Active principles with new modes of action should be well tolerated by crops, easy to apply, cost-effective, and amenable to regulation [98]. The agro-research industry is aiming to find effective herbicidal solutions to help sustain weed management diversity and crop production.

## 7. Conclusions and Recommendations on the Regulated Use of Glyphosate

The aim of the Farm to Fork strategy and the European Green Deal is to implement sustainable and environmentally friendly policies, in particular in agriculture, and this encompasses the protection of human health and biodiversity. Therefore, herbicides in general, and glyphosate in particular, are recovering our attention.

Glyphosate is a non-selective herbicide commonly used in croplands, urban areas, homes, and gardens. We can be exposed to this compound and its degradation components through the foods and the environment. In response to reclassification of glyphosate in Group 2A (probably carcinogenic to humans), regulations that are more stringent were implemented to set the maximum levels for glyphosate in livestock and poultry food products including meat, milk and eggs, but also in different crops.

Biological pest control, organic cropping, and regulative management help to reduce glyphosate use. Agro-ecology has drawn increasing interest and, according to many stakeholders, represents a strategic approach that can enable a successful transition to more sustainable farming and food systems. The policy in favour of agro-ecology would be exceptional, because it addresses all the levers needed to promote the agro-ecological transition, from production to consumption, by way of a transformation of the systems of education, research and development for achieving dietary security in the ever-lengthening shadow of climate change, bringing us thus closer to the realization of the plan in the 2030 Agenda for Sustainable Development and its 17 Sustainable Development Goals.

The potential carcinogenicity, massive use, and increasing presence of glyphosate residues in drinking water sources should lead regulatory agencies to take actions such as the following to protect human health: (a) Making trace level analyses in food and water mandatory; (b) re-assessing acceptable daily glyphosate intake levels; and (c) adding glyphosate to the water quality standards for drinking water sources. In this way, under the Sustainable Use of Pesticide Directives, each member state in the EU must have a national action plan that requires ‘quantitative objectives, targets, measures and timetables to reduce risks and impacts of pesticide use on human health and the environment’ which also includes measuring for residues. Secondly, they state a requirement for re-assessing the ADI values, but this already occurs when chemicals are re-evaluated to renew their licence to be on the European Market.

Environmental loads and exposure to glyphosate, AMPA, and formulation adjuvants continue to increase. Urban use of glyphosate in emerging countries is also a key issue. The European Food Safety Authority (EFSA) succeeded in identifying the potentially deleterious consequences of glyphosate on untargeted wild terrestrial vertebrates [118] in different scenarios such as crops pre-planting, post-planting and pre-emergence, cereals and oilseeds pre-harvest, and orchard crops and grapes.

There has been a strong controversy over the use of this herbicide and the detection of its residues in various foodstuffs. In 2013, the German Institute for Risk Assessment (BfR) conducted a comprehensive study concluding that classifying glyphosate as a carcinogen was unwarranted. This conclusion was reinforced by the EFSA in 2014 and 2015 by deeming it unlikely for glyphosate to pose serious risks to humans, in line with EPA’s statements of 1993, 2015, and 2016 that glyphosate was probably not carcinogenic. However, some of the scientific community refuted these claims, some agreed, and some just stated that

further research and work is required for this chemical, as it was already supported in previous sections. This is because no information about potential conflicts of interest of the authors of the reports was revealed or even that they were verbatim copies of previous studies produced by the multinational Monsanto, currently owned by Bayer. In that way, the public was served with controversy and serious doubts about potential prevarication on the part of the interested stakeholders. On top of that, in 2015, the WHO agency IARC classified glyphosate as probably carcinogenic to humans (Group 2A) after analysis of published scientific papers that associated to some cases of lymphomas (vide supra). These contradictions have led glyphosate use to being banned or strictly regulated in some countries. There were other issues with these evaluations, such as the IARC was evaluating for hazard, while the European Food Safety Authority (EFSA) and European Chemicals Agency (ECHA) were evaluating for risk. The difference between a hazard and risk assessment is the inclusion of probable exposure levels expected. Thus, the massive use of glyphosate and the expansion of glyphosate-resistant transgenic plants has allowed extensive agricultural production to be increased with substantially reduced costs. In this respect, the agrochemical can help fulfil the Millennium Goals and the 2030 Agenda by allowing adequate amounts of food to be supplied to a growing population. Moreover, it is clear that banning glyphosate use would increase tillage—and agricultural production—costs in some European Union countries. On a market with total price freedom, and under control of logistics and distribution multinationals, farmers are forced to keep prices low in order to compete with non-EU producers, many of whom can still use glyphosate virtually freely. As a result, EU farmers are having to sell their products at prices below their production costs, thus causing strong economic stress in European agriculture. In any case, there is a shadow of doubt about the safety of glyphosate as an herbicide, so appropriate measures should be taken in this respect to protect public health, and the environment. Furthermore, a need exists to avoid unfair competition from producers in countries where different regulatory values are in force.

It is key to finance epidemiological, biomonitoring, and toxicological research on endocrinology to verify if the consequences of glyphosate are mediated by endocrine disturbance, for example. That is the main conclusion that we want to make clear, recommending further follow-up to clarify this matter.

**Author Contributions:** Conceptualization, Y.S., J.-C.M., J.X. and J.S.-G.; resources, Y.S., J.-C.M., J.X. and J.S.-G.; writing—original draft preparation, Y.S., J.-C.M., J.X. and J.S.-G.; writing—review and editing, J.-C.M. and J.S.-G.; visualization, J.-C.M.; supervision, J.S.-G.; project administration, J.S.-G.; funding acquisition, J.S.-G. All authors have read and agreed to the published version of the manuscript.

**Funding:** Xunta de Galicia, ED431C2018/42-GRC.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Dill, G.M.; Sammons, R.D.; Feng, P.C.; Kohn, F.; Kretzmer, K.; Mehrsheikh, A.; Bleeke, M.; Honegger, J.L.; Farmer, D.; Wright, D.; et al. Glyphosate: Discovery, Development, Applications, and Properties. In *Glyphosate Resistance in Crops and Weeds: History, Development, and Management*; Nandula, V.K., Ed.; Wiley: Hoboken, NJ, USA, 2010; ISBN 978-0-470-41031-8.
2. Vandenberg, L.N.; Blumberg, B.; Antoniou, M.N.; Benbrook, C.M.; Carroll, L.; Colborn, T.; Everett, L.G.; Hansen, M.; Landrigan, P.J.; Lanphear, B.P.; et al. Is it time to reassess current safety standards for glyphosate-based herbicides? *J. Epidemiol. Commun. Health* **2017**, *71*, 613–618. [[CrossRef](#)] [[PubMed](#)]
3. Duke, S.O. Perspectives on transgenic, herbicide-resistant crops in the USA almost 20 years after introduction. *Pest Manag. Sci.* **2015**, *71*, 652–657. [[CrossRef](#)] [[PubMed](#)]
4. Duke, S.O.; Dayan, F.E. Discovery of new herbicide modes of action with natural phytotoxins. *Am. Chem. Soc. Symp. Ser.* **2015**, *1204*, 79–92. [[CrossRef](#)]

5. Benbrook, C.M. Impacts of genetically engineered crops on pesticide use in the U.S.—The first 16 years. *Environ. Sci. Eur.* **2012**, *24*, 24. [CrossRef]
6. Heap, I.M. Global perspective of herbicide-resistant weeds. *Pest Manag. Sci.* **2014**, *70*, 1306–1315. [CrossRef] [PubMed]
7. Mortensen, D.A.; Egan, J.F.; Maxwell, B.D.; Ryan, M.R. Navigating a critical juncture for sustainable weed management. *BioScience* **2012**, *62*, 75–84. [CrossRef]
8. NASS—National Agricultural Statistics Service. U.S. Soybean Industry: Glyphosate Effectiveness Declines. NASS Highlights N°2014-1. 2014. Available online: [https://www.nass.usda.gov/Surveys/Guide\\_to\\_NASS\\_Surveys/Ag\\_Resource\\_Management/ARMS\\_Soybeans\\_Factsheet/ARMS\\_2012\\_Soybeans.pdf](https://www.nass.usda.gov/Surveys/Guide_to_NASS_Surveys/Ag_Resource_Management/ARMS_Soybeans_Factsheet/ARMS_2012_Soybeans.pdf) (accessed on 14 June 2020).
9. Cerdeira, A.L.; Gazziero, D.L.P.; Duke, S.O.; Matallo, M.B. Agricultural impacts of glyphosate-resistant soybean cultivation in South America. *J. Agric. Food Chem.* **2011**, *59*, 5799–5807. [CrossRef]
10. Beckie, H.J. Herbicide-resistant weeds: Management tactics and practices. *Weed Technol.* **2006**, *20*, 793–814. [CrossRef]
11. Beckie, H.J.; Ashworth, M.B.; Flower, K.C. Herbicide Resistance Management: Recent Developments and Trends. *Plants* **2019**, *8*, 161. [CrossRef]
12. Moss, S.R. Integrated weed management (IWM): Why are farmers reluctant to adopt non-chemical alternatives to herbicides? *Pest Manag. Sci.* **2019**, *75*, 1205–1211. [CrossRef] [PubMed]
13. Giesy, J.P.; Dobson, S.; Solomon, K.R. Ecotoxicological risk assessment for Roundup® herbicide. *Rev. Environ. Cont. Toxicol.* **2000**, *167*, 35–120. [CrossRef]
14. Borggard, O.K.; Gimsing, A.L. Fate of glyphosate in soil and the possibility of leaching to ground and surface waters: A review. *Pest Manag. Sci.* **2008**, *64*, 441–456. [CrossRef] [PubMed]
15. Gillezeau, C.; Van Gerwen, M.; Shaffer, R.M.; Rana, I.; Zhang, L.; Sheppard, L.; Taioli, E. The evidence of human exposure to glyphosate: A review. *Environ. Health* **2019**, *18*, 2–14. [CrossRef] [PubMed]
16. Boobis, A.; Ossendorp, B.C.; Banasiak, U.; Hamey, P.Y.; Sebestyen, I.; Moretto, A. Cumulative risk assessment of pesticide residues in food. *Toxicol. Lett.* **2008**, *180*, 137–150. [CrossRef]
17. Maeda, H.; Dudareva, N. The shikimate pathway and aromatic amino acid biosynthesis in plants. *Ann. Rev. Plant Biol.* **2012**, *63*, 73–105. [CrossRef]
18. Heap, I. The International Herbicide-Resistant Weed Database. Available online: [www.weedscience.org](http://www.weedscience.org) (accessed on 7 December 2020).
19. FAO. *Management of Herbicide-Resistant Weed Populations: 100 Questions on Resistance*; FAO: Rome, Italy, 2008.
20. Beckie, H.J.; Heap, I.M.; Smeda, R.J.; Hall, I.M. Screening for herbicide resistance in weeds. *Weed Technol.* **2000**, *14*, 428–445. [CrossRef]
21. Gaines, T.A.; Ward, S.M.; Bukun, B.; Preston, C.; Leach, J.E.; Westra, P. Interspecific hybridization transfers a previously unknown glyphosate resistance mechanism in *Amaranthus* species. *Evol. Appl.* **2012**, *5*, 29–38. [CrossRef] [PubMed]
22. Ghanizadeh, H.; Harrington, K.C. Non-target site mechanisms of resistance to herbicides. *Crit. Rev. Plant Sci.* **2017**, *36*, 24–34. [CrossRef]
23. Jugulam, M.; Shyam, C. Non-Target-Site Resistance to herbicides: Recent developments. *Plants* **2019**, *8*, 417. [CrossRef]
24. Chueca, C.; Cirujeda, A.; De Prado, R.; Diaz, E.; Ortas, L.; Taberner, A.; Zaragoza, C. *Colección de Folletos Sobre Manejo de Poblaciones Resistentes en Papaver, Lolium, Avena y Echinochloa*; SEMh Grupo de Trabajo CPRH: Valencia, Spain, 2005.
25. Storrie, A. Herbicide Resistance Mechanisms and Common HR Misconceptions. *2006 Grains Research Update for Irrigation Croppers. Brochure. Switzerland.* 2006.
26. Moss, S.R. Herbicide resistance: New threats, new solutions? In Proceedings of the HGCA CONFERENCE. Arable Crop Protection in the Balance: Profit and the Environment, Grantham, UK, 25–26 January 2006.
27. Gage, K.L.; Krausz, R.F.; Walters, S.A. Emerging challenges for weed management in herbicide-resistant crops. *Agriculture* **2019**, *9*, 180. [CrossRef]
28. Reddy Krishna, N.; Jha, P. Herbicide-resistant weeds: Management strategies and upcoming technologies. *Indian J. Weed Sci.* **2016**, *48*, 108–111. [CrossRef]
29. Boerboom, C.; Owen, M. *Facts about Glyphosate-Resistant Weeds*; USDA: Washington, DC, USA, 2007.
30. Heap, I.; Duke, S.O. Overview of glyphosate-resistant weeds worldwide. *Pest Manag. Sci.* **2018**, *74*, 1040–1049. [CrossRef] [PubMed]
31. Owen, M.; Beckie, H.J.; Leeson, J.; Norsworthy, J.K.; Steckel, L.E. Integrated pest management and weed management. *Pest Manag. Sci.* **2014**, *71*, 357–376. [CrossRef] [PubMed]
32. Green, J.M. Current state of herbicides in herbicide-resistant crops. *Pest Manag. Sci.* **2014**, *70*, 1351–1357. [CrossRef] [PubMed]
33. Lombardo, L.; Coppola, G.; Zelasco, S. New Technologies for Insect-Resistant and Herbicide-Tolerant Plants. *Trends Biotechnol.* **2016**, *34*, 49–57. [CrossRef]
34. Gasser, C.S.; Fraley, R.T. Genetically Engineering Plants for Crop Improvement. *Science* **1989**, *244*, 1293–1299. [CrossRef]
35. Auer, C.; Frederick, R. Crop improvement using small RNAs: Applications and predictive ecological risk assessments. *Trends Biotechnol.* **2009**, *27*, 644–651. [CrossRef] [PubMed]
36. Runo, S.; Alakonya, A.; Machuka, J.; Sinha, N. RNA interference as a resistance mechanism against crop parasites in Africa: A ‘Trojan horse’ approach. *Pest Manag. Sci.* **2011**, *67*, 129–136. [CrossRef]
37. Green, J.M. The benefits of herbicide-resistant crops. *Pest Manag. Sci.* **2012**, *68*, 1323–1331. [CrossRef]

38. Espinoza, C.; Schelecheter, R.; Herrera, D.; Torres, E.; Serrano, A.; Medina, C.; Arce-Johnson, P. Cisgenesis and Intragenesis: New tools for Improving Crops. *Biol. Res.* **2013**, *46*, 323–331. [[CrossRef](#)]
39. Zhu, T.; Mettenburg, K.; Peterson, D.J.; Tagliani, L.; Baszczynski, C.L. Engineering herbicide-resistant maize using chimeric RNA/DNA oligonucleotides. *Nat. Biotechnol.* **2000**, *18*, 555–558. [[CrossRef](#)]
40. Sun, Y.; Zang, X.; Wu, C.; He, Y.; Ma, Y.; Hou, H.; Guo, X.; Du, W.; Zhaom, Y.; Xia, L. Engineering Herbicide-Resistant Rice Plants through CRISPR/Cas9-Mediated Homologous Recombination of Acetolactate Synthase. *Mol. Plant* **2016**, *9*, 628–631. [[CrossRef](#)]
41. Luo, M.; Gilbert, B.; Ayliffe, M. Applications of CRISPR/Cas9 technology for targeted mutagenesis, gene replacement and stacking of genes in higher plants. *Plant Cell Rep.* **2016**, *35*, 1439–1450. [[CrossRef](#)]
42. Han, Y.-J.; Kim, J.-I. Application of CRISPR/Cas9-mediated gene editing for the development of herbicide-resistant plants. *Plant Biotechnol. Rep.* **2019**, *13*, 447–457. [[CrossRef](#)]
43. Hussain, B.; Lucas, S.J.; Budak, H. CRISPR/Cas9 in plants: At play in the genome and at work for crop improvement. *Brief. Funct. Genom.* **2018**, *17*, 319–328. [[CrossRef](#)] [[PubMed](#)]
44. Tian, S.; Jiang, L.; Cui, X.; Zhang, J.; Guo, S.; Li, M.; Zhang, H.; Ren, Y.; Gong, G.; Zong, M.; et al. Engineering herbicide-resistant watermelon variety through CRISPR/Cas9-mediated base-editing. *Plant Cell Rep.* **2018**, *37*, 1353–1356. [[CrossRef](#)] [[PubMed](#)]
45. Zaidi, S.S.; Mahfouz, M.M.; Mansoor, S. CRISPR-Cpf1: A New Tool for Plant Genome Editing. *Trends Plant Sci.* **2017**, *22*, 550–553. [[CrossRef](#)] [[PubMed](#)]
46. Kamthan, A.; Chaudhuri, A.; Kamthan, M.; Datta, A. Genetically modified (GM) crops: Milestones and new advances in crop improvement. *Theor. Appl. Genet.* **2016**, *129*, 1639–1655. [[CrossRef](#)]
47. Soda, N.; Verma, L.; Giri, J. CRISPR-Cas9 based plant genome editing: Significance, opportunities and recent advances. *Plant Physiol. Biochem.* **2018**, *131*, 2–11. [[CrossRef](#)]
48. Ni, Z.; Han, Q.; He, Y.-Q.; Huang, S. Application of genome-editing technology in crop improvement. *Cereal Chem.* **2018**, *95*, 35–48. [[CrossRef](#)]
49. Weston, L.A.; Duke, S.O. Weed and Crop Allelopathy. *Crit. Rev. Plant Sci.* **2003**, *22*, 367–389. [[CrossRef](#)]
50. Iqbal, J.; Cheema, Z.A.; Mushtaq, M.N. Allelopathic Crop Water Extracts Reduce the Herbicide Dose for Weed Control in Cotton (*Gossypium hirsutum*). *Int. J. Agric. Biol.* **2009**, *11*, 360–366.
51. Fujii, Y. Screening and Future Exploitation of Allelopathic Plants as Alternative Herbicides with Special Reference to Hairy Vetch. *J. Crop Prod.* **2001**, *4*, 257–275. [[CrossRef](#)]
52. Belz, R.G. Allelopathy in crop/weed interactions—An update. *Pest Manag. Sci.* **2007**, *63*, 308–326. [[CrossRef](#)] [[PubMed](#)]
53. Shirgapure, K.H.; Ghosh, P. Allelopathy a Tool for Sustainable Weed Management. *Arch. Curr. Res. Int.* **2020**, *20*, 17–25. [[CrossRef](#)]
54. Farooq, N.; Abbs, T.; Tanveer, A.; Jabran, K. Allelopathy for Weed Management. In *Co-Evolution of Secondary Metabolites. Reference Series in Phytochemistry*; Mérillon, J.M., Ramawat, K., Eds.; Springer: Cham, Switzerland, 2020; ISBN 978-3-319-96396-9. [[CrossRef](#)]
55. Muhammad, Z.; Inayat, N.; Majeed, A.; Rehmanullah; Ali, H.; Ullah, K. Allelopathy and Agricultural Sustainability: Implication in weed management and crop protection—An overview. *Eur. J. Ecol.* **2019**, *5*, 54–61. [[CrossRef](#)]
56. Bajwa, A.A.; Khan, M.J.; Bhowmik, P.C.; Walsh, M.; Chauhan, B.S. Sustainable Weed Management. In *Innovations in Sustainable Agriculture*; Farooq, M., Pisante, M., Eds.; Springer: Cham, Switzerland, 2019; ISBN 978-3-030-23168-2. [[CrossRef](#)]
57. Kennedy, A.C. Soil Microorganisms for weed management. *J. Crops Prod.* **1999**, *2*, 123–138. [[CrossRef](#)]
58. Kennedy, A.C.; Kremer, R.J. Microorganisms in Weed Control Strategies. *J. Prod. Agric.* **1996**, *9*, 480–485. [[CrossRef](#)]
59. Kremer, R.J. Management of Weed Seed Banks with Microorganisms. *Ecol. Appl.* **1993**, *3*, 42–52. [[CrossRef](#)]
60. Boyetchko, S.M. Principles of Biological Weed Control with Microorganisms. *Hort. Sci.* **1997**, *32*, 201–205.
61. Kremer, R.J.; Caesar, A.J.; Souissi, T. Soilborne microorganisms of Euphorbia are potential biological control agents of the invasive weed leafy spurge. *Appl. Soil. Ecol.* **2006**, *32*, 27–37. [[CrossRef](#)]
62. Thrall, P.H.; Oakeshott, J.G.; Fitt, G.; Southerton, S.; Burdon, J.J.; Sheppard, A.; Russell, R.J. Evolution in agriculture: The application of evolutionary approaches to the management of biotic interactions in agro-ecosystems. *Evol. Appl.* **2011**, *4*, 200–215. [[CrossRef](#)]
63. Devine, M.D.; Shukla, A. Altered target sites as a mechanism of herbicide resistance. *Crop Protect.* **2000**, *19*, 881–889. [[CrossRef](#)]
64. Sibony, M.; Rubin, B. Molecular basis for multiple resistance to acetolactate synthase- inhibitin herbicides and atrazine in *Amarantus biotoides* (prostrate pigweed). *Planta* **2003**, *216*, 1022–1027. [[CrossRef](#)] [[PubMed](#)]
65. Li, J.; Smeda, R.J.; Nelson, K.A.; Dayan, F.E. Physiological basis for resistance to diphenyl ether herbicides in common waterhemp (*Amaranthus rudis*). *Weed Sci.* **2004**, *52*, 333–338. [[CrossRef](#)]
66. Neve, P. Challenges for herbicide resistance evolution and management: 50 years after Harper. *Weed Res.* **2007**, *47*, 365–369. [[CrossRef](#)]
67. Yu, Q.; Powles, S. Metabolism-Based Herbicide Resistance and Cross-Resistance in Crop Weeds: A Threat to Herbicide Sustainability and Global Crop Production. *Plant Physiol.* **2014**, *166*, 1106–1118. [[CrossRef](#)] [[PubMed](#)]
68. Busi, R.; Vila-Aiub, N.; Beckie, H.J.; Gaines, T.A.; Goggin, D.E.; Kaundun, S.S.; Lacoste, M.; Neve, P.; Nissen, S.J.; Norsworthy, J.K.; et al. Herbicide-resistant weeds: From research and knowledge to future needs. *Evol. Appl.* **2013**, *6*, 1218–1221. [[CrossRef](#)] [[PubMed](#)]
69. Busi, R.; Powles, S.B.; Beckie, H.J.; Renton, M. Rotations and mixtures of soil-applied herbicides delay resistance. *Pest Manag. Sci.* **2019**, *76*, 487–496. [[CrossRef](#)]



70. Vila-Aiub, M.M.; Gundel, P.; Yu, Q.; Powles, S.B. Glyphosate resistance in *Sorghum halepense* and *Lolium rigidum* is reduced at suboptimal growing temperatures. *Pest Manag. Sci.* **2013**, *69*, 228–232. [[CrossRef](#)] [[PubMed](#)]
71. Renton, M. Shifting focus from the population to the individual as a way forward in understanding, predicting and managing the complexities of evolution of resistance to pesticides. *Pest Manag. Sci.* **2013**, *69*, 171–175. [[CrossRef](#)]
72. Alonso, A.; Sánchez, P.; Martínez, J.L. Environmental selection of antibiotic resistance genes. *Environ. Microbiol.* **2001**, *3*, 1–9. [[CrossRef](#)]
73. Matzrafi, M.; Seiwert, B.; Reemtsma, T.; Rubin, B.; Peleg, Z. Climate change increases the risk of herbicide-resistant weeds due to enhanced detoxification. *Planta* **2016**, *244*, 1217–1227. [[CrossRef](#)]
74. Norsworthy, J.K.; Ward, S.M.; Shaw, D.R.; Llewellyn, R.S.; Nichols, R.L.; Webster, T.M.; Bradley, K.W.; Frisvold, G.; Powles, S.B.; Burgos, N.R.; et al. Reducing the risks of herbicide resistance: Best management practices and recommendations. *Weed Sci.* **2012**, *60*, 31–62. [[CrossRef](#)]
75. Perotti, V.E.; Larran, A.S.; Palmieri, V.E.; Martinatto, A.K.; Permingeat, H.R. Herbicide resistant weeds: A call to integrate conventional agricultural practices, molecular biology knowledge and new technologies. *Plant Sci.* **2020**, *290*, 110255. [[CrossRef](#)] [[PubMed](#)]
76. Shaner, D.N. Lessons learned from the history of herbicide resistance. *Weed Sci.* **2014**, *62*, 427–431. [[CrossRef](#)]
77. Rosset, J.D.; Gulden, R.H. Cultural weed management practices shorten the critical weed-free period for soybean grown in the Northern Great Plains. *Weed Sci.* **2020**, *68*, 79–91. [[CrossRef](#)]
78. Duary, B. Weed prevention for quality seed production of crops. *SATSA Mukhapatra Annu. Tech. Issue* **2014**, *18*, 48–57.
79. IAASTD. *Agriculture at a Crossroads. International Assessment of Agricultural Knowledge, Science and Technology for Development*; Synthesis Report; IAASTD: Washington, DC, USA, 2008.
80. Gaba, S.; Fried, G.; Kazakou, E.; Chauvel, B.; Navas, M.-L. Agroecological weed control using a functional approach: A review of cropping systems diversity. *Agric. Sustain. Dev.* **2013**, *34*, 1–17. [[CrossRef](#)]
81. Macías, F.A. Allelopathy in the research of natural herbicide models. *ACS Symp. Ser.* **1995**, *582*, 310–329. [[CrossRef](#)]
82. Kelton, J.; Price, A.J.; Mosjidis, J. Allelopathic weed suppression through the use of cover crops. In *Weed Control*; Price, A.J., Ed.; Intech Press: Rijeka, Croatia, 2012; pp. 115–130. ISBN 978-953-51-0159-8/978-953-51-5215-6. [[CrossRef](#)]
83. Isman, M.B. Botanical insecticides, deterrents and repellents in modern agriculture and an increasingly regulated world. *Ann. Rev. Entomol.* **2006**, *51*, 45–66. [[CrossRef](#)] [[PubMed](#)]
84. Macías, F.A.; Molinillo, J.M.G.; Varela, R.M.; Galindo, J.C.G. Allelopathy, a natural alternative for weed control. *Pest Manag. Sci.* **2007**, *63*, 327–384. [[CrossRef](#)] [[PubMed](#)]
85. Tabaglio, V.; Gavazzi, C.; Schulz, M.; Marocco, A. Alternative weed control using the allelopathic effect of natural bezoxazinoids from rye mulch. *Agron. Sustain. Dev.* **2008**, *28*, 397–401. [[CrossRef](#)]
86. Khalid, S.; Ahmand, T.; Shad, R.A. Use of allelopathy in agriculture. *Asian J. Plant Sci.* **2002**, *1*, 292–297. [[CrossRef](#)]
87. Vyvyan, J.R. Allelochemicals as leads for new herbicides and agrochemicals. *Tetrahedron* **2002**, *58*, 1631–1646. [[CrossRef](#)]
88. Soltys, D.; Krasuska, U.; Bogatek, R.; Gniazdowska, A. Allelochemicals as bioherbicides: Present and perspectives. In *Herbicides: Current Research and Cases Studies in Use*; Price, A., Kelton, J., Eds.; IntechOpen: London, UK, 2013; ISBN 978-953-51-1112-2. [[CrossRef](#)]
89. Balah, M.A. Formulation of prospective plant oils derived micro-emulsions for herbicidal activity. *J. Plant Prot. Path.* **2013**, *4*, 911–926.
90. Wang, Y.; Zhao, R.; Yu, L.; Zhang, Y.; He, Y.; Yao, J. Evaluation of cinnamon essential oil microemulsion and its vapor phase for controlling postharvest gray mold of pears (*Pyrus pyrifolia*). *J. Sci. Food Agric.* **2014**, *30*, 1000–1004. [[CrossRef](#)]
91. Massoud, A.; Manal, M.A.; Osman, A.Z.; Magdy, I.E.M.; Abdel-Rheim, K.H. Eco-Friendly Nano-emulsion Formulation of *Mentha piperita* Against Stored Product Pest *Sitophilus oryzae* Magdy. *Adv. Crop Sci. Tech.* **2018**, *6*, 1000404. [[CrossRef](#)]
92. Rakmai, J.; Cheirsilp, B.; Cid, A.; Torrado-Agrasar, A.; Mejuto, J.C.; Simal-Gandara, J. Encapsulation of Essential Oils by Cyclodextrins: Characterization and Evaluation. In *Cyclodextrin: A Versatile Ingredient*; Arora, P., Dhingra, N., Eds.; IntechOpen: London, UK, 2018; pp. 264–290. ISBN 978-1-78923-068-0/978-1-83881-379-6. [[CrossRef](#)]
93. Rakmai, J.; Cheirsilp, B.; Mejuto, J.C.; Simal-Gándara, J.; Torrado-Agrasar, A. Antioxidant and antimicrobial properties of encapsulated guava leaf oil in hydroxypropyl-beta-cyclodextrin. *Ind. Crops Prod.* **2018**, *111*, 219–225. [[CrossRef](#)]
94. Rakmai, J.; Cheirsilp, B.; Cid, A.; Torrado-Agrasar, A.; Simal-Gandara, J.; Mejuto, J.C. Encapsulation of yarrow essential oil in hydroxypropyl-beta-cyclodextrin: Physicochemical characterization and evaluation of bio-efficacies. *CyTA J. Food.* **2017**, *15*, 409–417. [[CrossRef](#)]
95. Rakmai, J.; Cheirsilp, B.; Mejuto, J.C.; Torrado-Agrasar, A.; Simal-Gandara, J. Physico-chemical characterization and evaluation of bio-efficacies of black pepper essential oil encapsulated in hydroxypropyl-beta-cyclodextrin. *Food Hydrocol.* **2017**, *65*, 157–164. [[CrossRef](#)]
96. Webber III, C.L.; Shrefler, J.W.; Brandenberger, L.P. Organic Weed Control. In *Herbicides: Environmental Impact Studies and Management Approaches*; Alvarez-Fernandez, R., Ed.; Intech Open: London, UK, 2012; pp. 185–189. ISBN 978-953-307-892-2/978-953-51-5181-4. [[CrossRef](#)]
97. Singh, N.; Chaput, L.; Villoutreix, B.O. Virtual screening web servers: Designing chemical probes and drug candidates in the cyberspace. *Brief. Bioinform.* **2020**, bbaa034. [[CrossRef](#)] [[PubMed](#)]

98. Kraehmer, H.; Laber, B.; Rosinger, C.; Schulz, A. Herbicides as weed control agents: State of the art: I. Weed control research and safener technology: The path to modern agriculture. *Plant Physiol.* **2014**, *166*, 1119–1131. [[CrossRef](#)]
99. Angeline, L.G.; Carpanese, G.; Cioni, P.L.; Morelli, I.; Macchia, M.; Flamini, G. Essential oils from Mediterranean *Lamiaceae* as weed germination inhibitors. *J. Agric. Food Chem.* **2003**, *51*, 6158–6164. [[CrossRef](#)] [[PubMed](#)]
100. Tworokski, T. Herbicide effects of essential oils. *Weed Sci.* **2002**, *50*, 425–431. [[CrossRef](#)]
101. Dayan, F.E.; Cantrell, C.L.; Duke, S.O. Natural products in crop protection. *Bioorg. Med. Chem.* **2009**, *17*, 4022–4034, and references therein. [[CrossRef](#)]
102. Santana, O.; Cabrera, R.; Giménez, C.; González-Coloma, A.; Sánchez-Vioque, R.; de los Mozos-Pascual, M.; Rodríguez-Conde, M.F.; Laserna-Ruiz, I.; Usano-Aleman, J.; Herraiz, D. Chemical and biological profiles of the essential oils from aromatic plants of agro industrial interest in Castilla-La Mancha (Spain). *Grasas Aceites Int. J. Fats Oils* **2012**, *63*, 214–222. [[CrossRef](#)]
103. Mucciarelli, M.; Camusso, W.; Bertea, C.M.; Maffei, C.M. Effect of (+)-pulegone and other oil components of *Mentha piperita* on cucumber respiration. *Phytochemistry* **2001**, *57*, 91–98. [[CrossRef](#)]
104. Verdeguer, M.; Catañeda, L.G.; Torre-Pagan, N.; Llorens-Molina, J.A.; Carrubba, A. Control of *Erierson bonariensis* with *Thymbra captata*, *Mentha piperitum* *Eucalyptus camaldulensis* and *Santolina chamaecyparissus* essential oils. *Molecules* **2020**, *25*, 562. [[CrossRef](#)] [[PubMed](#)]
105. Vasilakoglou, I.; Dhima, K.; Wogiatzi, E.; Eleftherohorinos, I.; Lithourgidis, A. Herbicidal potential of essential oils of oregano or marjoram (*Origanum* ssp.) and Basil (*Ocimum basilicum*) on *Echinochloa crus-galli* (L.) P. Beauv. and *Chenopodium album* L. weeds. *Allelopat. J.* **2007**, *20*, 297–306.
106. García-Rellán, D.; Verdeguer, M.; Salamone, A.; Blázquez, M.A.; Boira, H. Chemical composition, herbicidal and antifungal activity of *Satureja cuneifolia* essential oils from Spain. *Nat. Prod. Commun.* **2016**, *11*, 841–844. [[CrossRef](#)] [[PubMed](#)]
107. Zhang, J.; An, M.; Wu, H.; Liu, L.L.; Stanton, R. Chemical composition of essential oils of four *Eucalyptus* species and their phytotoxicity on silverleaf nightshade (*Solanum elaeagnifolium* Cav.) in Australia. *Plant Growth Reg.* **2012**, *68*, 231–237. [[CrossRef](#)]
108. Maaloul, A.; Verdeguer-Sancho, M.M.; Oddo, M.; Saadaoui, E.; Jebri, M.; Michalet, S.; Dijoux-Franca, M.G.; Mars, M.; Romdhane, M. Effect of Short and Long Term Irrigation with Treated Wastewater on Chemical Composition and Herbicidal Activity of *Eucalyptus camaldulensis* Dehn. Essential Oils. *Not. Bot. Horti Agrobot. Cluj Napoca* **2019**, *47*, 1374–1381. [[CrossRef](#)]
109. Barbosa, J.C.A.; Filomen, C.A.; Teixeira, R.R. Chemical variability and biological activities of *Eucalyptus* spp. Essential oils. *Molecules* **2016**, *21*, 1671. [[CrossRef](#)]
110. Ben Ghnaya, A.; Hamrouni, L.; Ahouses, I.; Hanana, M.; Romane, A. Study of allelopathic effects of *Eucalyptus erythrocorys* L. crude extracts against germination and seedling growth of weeds and wheat. *Nat. Prod. Res.* **2016**, *30*, 2058–2064. [[CrossRef](#)]
111. Ismail, A.; Lamia, H.; Mohsen, H.; Bassem, J. Herbicidal potential of essential oils from three Mediterranean trees on different weeds. *Curr. Bioact. Comp.* **2013**, *8*, 3–12. [[CrossRef](#)]
112. Verdeguer, M.; Blázquez, M.A.; Boira, H. Chemical composition and herbicidal activity of the essential oil from a *Cistus ladanifer* population from Spain. *Nat. Prod. Res.* **2012**, *26*, 1602–1609. [[CrossRef](#)]
113. Davoren, M.J.; Schiestl, R.H. Glyphosate based herbicides and cancer risk: A post IARC decision review of potential mechanisms, policy, and avenues of research. *Carcinogenesis* **2018**, *39*, 1207–1215. [[CrossRef](#)]
114. Sviridov, A.V.; Shushkova, T.V.; Ermakova, I.T.; Ivanova, E.V.; Epiktetov, D.O.; Leontievsky, A.A. Microbial degradation of glyphosate herbicides. *Appl. Biochem. Microbiol.* **2015**, *51*, 188–195. [[CrossRef](#)]
115. Rodríguez, J.; Perez, B.; Nebot, C.; Falque, E.; Simal-Gandara, J. Food production link to underground waters quality in A Limia river basin. *Agric. Ecosys. Environ.* **2020**, *297*, 106969. [[CrossRef](#)]
116. Singh, B.; Singh, K. Microbial degradation of herbicides. *Crit. Rev. Microbiol.* **2016**, *42*, 245–261. [[CrossRef](#)]
117. Van Bruggen, A.H.C.; He, M.M.; Shin, K.; Mai, V.; Jeong, K.C.; Finckh, M.R.; Morris, J.G. Environmental and health effects of the herbicide glyphosate. *Sci. Total Environ.* **2018**, *616*, 255–268. [[CrossRef](#)] [[PubMed](#)]
118. EFSA. Conclusion on the peer review of the pesticide risk assessment of the active substance glyphosate: Peer review of the pesticide risk assessment of the active substance glyphosate. *EFSA J.* **2015**, *13*, 4302. [[CrossRef](#)]
119. Hanna, D.E.L.; Tomscha, S.A.; Dallaire, C.O.; Bennett, E.M. A review of riverine ecosystem service quantification: Research gaps and recommendations. *J. Appl. Ecol.* **2017**, *55*, 1299–1311. [[CrossRef](#)]
120. Battin, T.J.; Besemer, K.; Bengtsson, M.M.; Romani, A.M.; Packmann, A.I. The ecology and biogeochemistry of stream biofilms. *Nat. Rev. Microbiol.* **2016**, *14*, 251–263. [[CrossRef](#)] [[PubMed](#)]
121. Vera, M.S.; Lagomarsino, L.; Sylvester, M.; Perez, G.L.; Rodriguez, P.; Mugni, H.; Sinistro, R.; Ferraro, M.; Bonetto, C.; Zagarese, H.; et al. New evidences of Roundup® (glyphosate formulation) impact on the periphyton community and the water quality of freshwater ecosystems. *Ecotoxicology* **2010**, *19*, 710–721. [[CrossRef](#)] [[PubMed](#)]
122. Lasier, P.J.; Urich, M.L.; Hassan, S.M.; Jacobs, W.N.; Bringolf, R.B.; Owens, K.M. Changing agricultural practices: Potential consequences to aquatic organisms. *Environ. Monit. Assess.* **2016**, *188*, 672. [[CrossRef](#)]
123. Hove-Jensen, B.; Zechel, D.L.; Jochimsen, B. Utilization of glyphosate as phosphate source: Biochemistry and genetics of bacterial carbon-phosphorus lyase. *Microbiol. Mol. Biol. Rev.* **2014**, *78*, 176–197. [[CrossRef](#)]
124. Krzysko-Lupicka, T.; Krecidlo, L.; Koszalkowska, M. The ability of selected bacteria to grow in the presence of glyphosate. *Ecol. Chem. Eng. Chem. Inzynieria Ekol. A* **2015**, *22*, 185–193. [[CrossRef](#)]
125. Klátyik, S.; Takács, E.; Mörtl, M.; Földi, A.; Trábert, Z.; Ács, É.; Darvas, B.; Székács, A. Dissipation of the herbicide active ingredient glyphosate in natural water samples in the presence of biofilms. *Int. J. Environ. Anal. Chem.* **2017**, *97*, 901–921. [[CrossRef](#)]

126. Grandcoin, A.; Piel, S.; Baures, E. Aminomethylphosphonic acid (AMPA) in natural waters: Its sources, behavior and environmental fate. *Water Res.* **2017**, *117*, 187–197. [[CrossRef](#)]
127. Carles, L.; Gardon, H.; Joseph, L.; Sanchis, J.; Farre, M.; Artigas, J. Meta-analysis of glyphosate contamination in surface waters and dissipation by biofilms. *Environ. Intern.* **2019**, *124*, 284–293. [[CrossRef](#)]
128. Battaglin, W.A.; Meyer, M.T.; Kuivila, K.M.; Dietze, J.E. Glyphosate and its degradation product AMPA occur frequently and widely in US soils, surface water, groundwater, and precipitation. *J. Am. Water Resour. Assoc.* **2014**, *50*, 275–290. [[CrossRef](#)]
129. Struger, J.; Van Stempvoort, D.R.; Brown, S.J. Sources of aminomethylphosphonic acid (AMPA) in urban and rural catchments in Ontario, Canada: Glyphosate or phosphonates in wastewater? *Environ. Pollut.* **2015**, *204*, 289–297. [[CrossRef](#)]
130. Heber, M.P.; Fugere, V.; Gonzalez, A. The overlooked impact of rising glyphosate use on phosphorus loading in agricultural watersheds. *Front. Ecol. Environ.* **2019**, *17*, 48–56. [[CrossRef](#)]
131. Aparicio, V.C.; De Gerónimo, E.; Marino, D.; Primost, J.; Carriquiriborde, P.; Costa, J.L. Environmental fate of glyphosate and aminomethylphosphonic acid in surface waters and soil of agricultural basins. *Chemosphere* **2013**, *93*, 1866–1873. [[CrossRef](#)] [[PubMed](#)]
132. Coupe, R.H.; Kalkhoff, S.J.; Capel, P.D.; Gregoire, C. Fate and transport of glyphosate and aminomethylphosphonic acid in surface waters of agricultural basins. *Pest Manag. Sci.* **2011**, *68*, 16–30. [[CrossRef](#)]
133. Poiger, T.; Buerge, I.J.; Bächli, A.; Müller, M.D.; Balmer, M.E. Occurrence of the herbicide glyphosate and its metabolite AMPA in surface waters in Switzerland determined with on-line solid phase extraction LC-MS/MS. *Environ. Sci. Poll. Res.* **2016**, *24*, 1588–1596. [[CrossRef](#)] [[PubMed](#)]
134. Fiorino, E.; Sehonova, P.; Plhalova, L.; Blahova, J.; Svobodova, Z.; Faggio, C. Effects of glyphosate on early life stages: Comparison between *Cyprinus carpio* and *Danio rerio*. *Environ. Sci. Poll. Res.* **2018**, *25*, 8542–8549. [[CrossRef](#)] [[PubMed](#)]
135. Mercurio, P.; Flores, F.; Mueller, J.F.; Carter, S.; Negri, A.P. Glyphosate persistence in seawater. *Mar. Poll. Bull.* **2014**, *85*, 385–390. [[CrossRef](#)]
136. Toss, V.; Leito, I.; Yurchenko, S.; Freiberg, R.; Kruve, A. Determination of glyphosate in surface water with high organic matter content. *Environ. Sci. Poll. Res.* **2017**, *24*, 7880–7888. [[CrossRef](#)]
137. Pizarro, H.; Vera, M.S.; Vinocur, A.; Pérez, G.; Ferraro, G.; Menéndez-Helman, R.J.; Dos Santos, A.M. Glyphosate input modifies microbial community structure in clear and turbid freshwater systems. *Environ. Sci. Poll.* **2015**, *23*, 5143–5153. [[CrossRef](#)]
138. Solomon, K.R. Estimated exposure to glyphosate in humans via environmental, occupational, and dietary pathways: An updated review of the scientific literature. *Pest. Manag. Sci.* **2019**, *76*, 2878–2885. [[CrossRef](#)] [[PubMed](#)]
139. Bonansea, R.I.; Filippi, I.; Wunderlin, D.A.; Marino, D.J.G.; Ame, M.V. The fate of glyphosate and AMPA in a freshwater endorheic basin: An ecotoxicological risk assessment. *Toxics* **2018**, *6*, 3. [[CrossRef](#)] [[PubMed](#)]
140. Annett, R.; Hamid, R.H.; Hontela, A. Impact of glyphosate and glyphosate-based herbicides on the freshwater environment. *J. Appl. Toxicol.* **2014**, *34*, 458–479. [[CrossRef](#)]
141. Thompson, D.G.; Wojtaszek, B.F.; Staznik, B.; Chartrand, D.T.; Stephenson, G.R. Chemical and biomonitoring to assess potential acute effects of vision herbicide on native amphibian larvae in forest wetlands. *Environ. Toxicol. Chem.* **2004**, *23*, 843–849. [[CrossRef](#)]
142. Matozzo, V.; Munari, M.; Masiero, L.; Finos, L.; Gabriella, M. Ecotoxicological hazard of a mixture of glyphosate and aminomethylphosphonic acid to the mussel *Mytilus galloprovincialis* (Lamarck 1819). *Sci. Rep.* **2019**, *9*, 14302. [[CrossRef](#)]
143. Refatti, J.P.; de Avila, L.A.; Noldin, J.A.; Pacheco, I.; Ribeiro-Pestana, R. Leaching and residual activity of imidazolinone herbicides in lowland soils. *Ciênc. Rural* **2017**, *47*, 1–6. [[CrossRef](#)]
144. Bermudez-Couso, A.; Arias-Estevéz, M.; Novoa-Muñoz, J.C.; Lopez-Periago, E.; Soto-Gonzalez, B.; Simal-Gandara, J. Seasonal distributions of fungicides in soils and sediments of a small river basin partially devoted to vineyards. *Water Res.* **2007**, *41*, 4515–4525. [[CrossRef](#)] [[PubMed](#)]
145. EPA. *Appendix to Bioaccumulation Testing and Interpretation for the Purpose of Sediment Quality Assessment: Status and Needs: Chemicals-Specific Summary Tables*; EPA-823-R-00-002; USEPA Office of Water: Washington, DC, USA, 2000.
146. Stewardson, M.J.; Shang, W.; Kattel, G.R.; Webb, J.A. Chapter 22—Environmental Water and Integrated Catchment Management. In *Water for the Environment from Policy and Science to Implementation and Management*; Horne, A.C., Webb, J.A., Stewardson, M.J., Richter, B., Acremanm, M., Eds.; Academic Press: Cambridge, MA, USA, 2017; pp. 519–536. [[CrossRef](#)]
147. Vymazal, J.; Brezinova, T. The use of constructed wetlands for removal of pesticides from agricultural runoff and drainage: A review. *Environ. Int.* **2015**, *75*, 11–20. [[CrossRef](#)]
148. Akingbemi, B.T.; Sottas, C.M.; Loulova, A.I.; Klonefelter, G.; Hardy, M.P. Inhibition of testicular steroidogenesis by the xenoestrogen bisphenol A is associated with reduced pituitary luteinizing hormone secretion and decreased steroidogenic enzyme gene expression in rat Leydig cells. *Endocrinology* **2004**, *145*, 592–603. [[CrossRef](#)] [[PubMed](#)]
149. Relyea, R.A. New effects of Roundup on amphibians: Predators reduce herbicide mortality; herbicides induce antipredator morphology. *Ecol. Appl.* **2012**, *22*, 634–647. [[CrossRef](#)]
150. Dimitrov, B.D.; Gadeva, P.G.; Benova, D.K.; Bineva, M.V. Comparative genotoxicity of the herbicides Roundup, Stomp and Reglone in plant and mammalian test systems. *Mutagenesis* **2006**, *21*, 375–382. [[CrossRef](#)] [[PubMed](#)]
151. Dallegrave, E.; Mantese, F.D.; Coelho, R.S.; Pereira, J.D.; Dalsenter, P.R.; Langeloh, A. The teratogenic potential of the herbicide Glyphosate-Roundup in Wistar rats. *Toxicol. Lett.* **2003**, *142*, 45–52. [[CrossRef](#)]

152. Mladinic, M.; Berend, S.; Vrdoljak, A.L.; Kopjar, N.; Radic, B.; Zeljezic, D. Evaluation of genome damage and its relation to oxidative stress induced by glyphosate in human lymphocytes in vitro. *Environ. Mol. Mutagen.* **2009**, *50*, 800–807. [[CrossRef](#)]
153. Wang, Y.; Ezemaduka, A.N.; Li, Z.; Chen, Z.; Song, C. Joint Toxicity of Arsenic, Copper and Glyphosate on Behavior, Reproduction and Heat Shock Protein Response in *Caenorhabditis elegans*. *Bull. Environ. Cont. Toxicol.* **2017**, *98*, 465–471. [[CrossRef](#)] [[PubMed](#)]
154. Mladinic, M.; Perkovic, P.; Zeljezic, D. Characterization of chromatin instabilities induced by glyphosate, terbuthylazine and carbofuran using cytome FISH assay. *Toxicol. Lett.* **2009**, *189*, 130–137. [[CrossRef](#)] [[PubMed](#)]
155. García-Pérez, J.A.; Alarcón-Gutiérrez, E.; Perroni, Y.; Barois, I. Earthworm communities and soil properties in shaded coffee plantations with and without applications of glyphosate. *Appl. Soil Ecol.* **2014**, *83*, 230–237. [[CrossRef](#)]
156. Mercado, S.A.S.; Quintero-Caleño, J.D. Cytotoxic evaluation of glyphosate, using *Allium cepa* L. as bioindicator. *Sci. Total Environ.* **2020**, *700*, 134452. [[CrossRef](#)] [[PubMed](#)]
157. Da Cruz, C.; Carraschi, S.P.; Shoigiri, N.S.; da Silva, A.F.; Pitelli, R.A.; Fernandes-Machado, M.R. Sensitivity, ecotoxicity and histopathological effects on neotropical fish exposed to glyphosate alone and associated to surfactant. *J. Environ. Chem. Ecotoxicol.* **2016**, *8*, 25–33. [[CrossRef](#)]
158. Beltrano, J.; Ruscitti, M.; Arago, C.; Ronco, M. Changes in the accumulation of shikimic acid in mycorrhized *Capsicum annuum* L. grown with application of glyphosate and phosphorus. *Theor. Exp. Plant Physiol.* **2013**, *25*, 2. [[CrossRef](#)]
159. Qin, Y.; Li, X.; Xiang, Y.; Wu, D.; Bai, L.; Li, Z.; Liang, Y. Toxic effects of glyphosate on diploid and triploid fin cell lines from *Misgurnus anguillicaudatus*. *Chemosphere* **2017**, *180*, 356–364. [[CrossRef](#)]
160. Batista, P.F.; Costa, A.C.; Megguer, C.A.; Lima, J.S.; Guimaraes, D.S.; Almeida, G.M.; Nascimento, K.J.T. Pouteria torta: A native species of the Brazilian Cerrado as a bioindicator of glyphosate action. *Braz. J. Biol.* **2018**, *78*, 2. [[CrossRef](#)] [[PubMed](#)]
161. Mottier, A.; Kientz-Bouchart, V.; Serpentine, A.; Lebel, J.M.; Jha, A.N.; Costil, K. Effects of glyphosate-based herbicides on embryo-larval development and metamorphosis in the Pacific oyster, *Crassostrea gigas*. *Aquat. Toxicol.* **2013**, *128–129*, 67–78. [[CrossRef](#)] [[PubMed](#)]
162. Mona, M.H.; Gaafar, R.M.; Helal, I.B.; Omran, N.E.; Slama, W.M. Evaluation of cytotoxic effects of atrazine and glyphosate herbicides on *Biomphalaria glabrata* snails. *J. Basic Appl. Zool.* **2013**, *66*, 68–75. [[CrossRef](#)]
163. EPA. *Glyphosate. Draft Human Health Risk Assessment in Support of Registration Review*; D417700. EPA-HQ-OPP-2009-0361-0068; Office of Pesticide Programs: Washington, DC, USA, 2017.
164. EPA. Revised Glyphosate Issue Paper: Evaluation of Carcinogenic Potential. 2017. Available online: [https://cfpub.epa.gov/si/si\\_public\\_file\\_download.cfm?p\\_download\\_id=534487](https://cfpub.epa.gov/si/si_public_file_download.cfm?p_download_id=534487) (accessed on 14 June 2020).
165. Guyton, K.Z.; Loomis, D.; Grosse, Y.; Ghissassi, F.E.; Benbrahim-Tallaa, L.; Guha, N.; Scoccianti, C.; Mattock, H.; Straif, K. Carcinogenicity of tetrachlorvinphos, parathion, malathion, diazinon, and glyphosate. *Lancet Oncol.* **2015**, *16*, 490–491. [[CrossRef](#)]
166. Karunanayake, C.P.; Spinelli, J.J.; McLaughlin, J.R.; Dosman, J.A.; Pahwa, P.; McDuffie, H.H. Hodgkin lymphoma and pesticides exposure in men: A Canadian case-control study. *J. Agromed.* **2012**, *17*, 30–39. [[CrossRef](#)] [[PubMed](#)]
167. Schinasi, L.; Leon, M.E. Non-Hodgkin lymphoma and occupational exposure to agricultural pesticide chemical groups and active ingredients: A systematic review and meta-analysis. *Int. J. Environ. Res. Public Health* **2014**, *11*, 4449–4527. [[CrossRef](#)] [[PubMed](#)]
168. De Roos, A.J.; Blair, A.; Rusiecki, J.; Hoppin, J.A.; Svec, M.; Dosemeci, M.; Sandler, D.P.; Alavanja, M.C. Cancer incidence among glyphosate-exposed pesticide applicators in the Agricultural Health Study cohort. *Environ. Health Perspect.* **2005**, *113*, 49–54. [[CrossRef](#)]
169. De Roos, A.J.; Blair, A.; Rusiecki, J.; Hoppin, J.A.; Svec, M.; Dosemeci, M.; Sandler, D.P.; Alavanja, M.C. Glyphosate Results Revisited. De Roos et al. Respond. *Environ. Health Perspect.* **2005**, *113*, A366–A367. [[CrossRef](#)]
170. Andreotti, G.; Koutros, S.; Hofmann, J.N.; Sandler, D.P.; Lubin, J.H.; Lynch, C.F.; Lerro, C.C.; De Roos, A.J.; Parks, C.G.; Alavanja, M.C.; et al. Glyphosate use and cancer incidence in the agricultural health study. *J. Natl. Cancer Inst.* **2018**, *110*, 509–516. [[CrossRef](#)] [[PubMed](#)]
171. Marc, J.; Mulner-Lorillon, O.; Boulben, S.; Hureau, D.; Durand, G.; Bellé, R. Pesticide Roundup provokes cell division dysfunction at the level of CDK1/cyclin B activation. *Chem. Res. Toxicol.* **2002**, *15*, 326–331. [[CrossRef](#)]
172. Marc, J.; Mulner-Lorillon, O.; Bellé, R. Glyphosate-based pesticides affect cell cycle regulation. *Biol. Cell.* **2004**, *96*, 245–249. [[CrossRef](#)]
173. Levine, S.L.; Han, Z.; Farmer, D.R.; Padopoulos, V. Disrupting mitochondrial function with surfactants inhibits MA-10 Leydig cell steroidogenesis. *Cell Biol. Toxicol.* **2007**, *23*, 385–400. [[CrossRef](#)] [[PubMed](#)]
174. Mesnage, R.; Bernay, B.; Seralini, G.E. Ethoxylated adjuvants of glyphosate-based herbicides are active principles of human cell toxicity. *Toxicology* **2013**, *313*, 122–128. [[CrossRef](#)] [[PubMed](#)]
175. Richard, S.; Moslemi, S.; Sipahutar, H.; Benachour, N.; Seralini, G.E. Differential effects of glyphosate and roundup on human placental cells and aromatase. *Environ. Health Perspect.* **2005**, *113*, 716–720. [[CrossRef](#)] [[PubMed](#)]
176. Guilherme, S.; Gaivao, I.; Santos, M.A.; Pacheco, M. European eel (*Anguilla anguilla*) genotoxic and pro-oxidant responses following short-term exposure to Roundup—A glyphosate-based herbicide. *Mutagenesis* **2010**, *25*, 523–530. [[CrossRef](#)]
177. Benbrook, C.M. How did the US EPA and IARC reach diametrically opposed conclusions on the genotoxicity of glyphosate-based herbicides? *Environ. Sci. Eur.* **2019**, *31*, 2. [[CrossRef](#)]
178. Martens, M.A.; Bleeke, M.S.; Leopold, V.A.; Farmer, D.R. Toxicology and human health risk assessment of polyethoxylated tallow amine surfactant used in glyphosate formulations. *Regul. Toxicol. Pharmacol.* **2019**, *107*, 104347. [[CrossRef](#)]

179. IARC. *Glyphosate Monograph*; International Agency for Research on Cancer, World Health Organization: Geneva, Switzerland, 2015; Volume 112.
180. Leon, M.E.; Schinasi, L.H.; Lebailly, P.; Freeman, L.E.B.; Nordby, K.C.; Ferro, G.; Monnereau, A.; Brouwer, M.; Tual, S.; Baldi, I.; et al. Pesticide use and risk of non-Hodgkin lymphoid malignancies in agricultural cohorts from France, Norway and the USA: A pooled analysis from the AGRICOH consortium. *Int. J. Epidemiol.* **2019**, *48*, 1519–1535. [CrossRef] [PubMed]
181. Zhang, L.; Rana, I.; Taioli, E.; Shaffer, R.M.; Sheppard, L. Exposure to Glyphosate-Based Herbicides and Risk for Non-Hodgkin Lymphoma: A Meta-Analysis and Supporting Evidence. *Mutat. Res.* **2019**, *781*, 186–206. [CrossRef]
182. Amrhein, V.; Greenland, S.; McShane, B. Scientists rise up against statistical significance. *Nature* **2019**, *567*, 305–307. [CrossRef] [PubMed]
183. EPA. Email from Vincent Cogliano to Norman Birchfield, both of EPA's Office of Research and Development (ORD), 12/7/15. 2015. Available online: <https://assets.documentcloud.org/documents/4641115/Cogliano-Memo.pdf> (accessed on 15 June 2020).
184. Romano, R.M.; Romano, M.A.; Bernardi, M.M.; Furtado, P.V.; Oliveira, C.A. Prepubertal exposure to commercial formulation of the herbicide glyphosate alters testosterone levels and testicular morphology. *Arch. Toxicol.* **2010**, *84*, 309–317. [CrossRef]
185. Gasnier, C.; Dumont, C.; Benachour, N.; Clair, E.; Chagnon, M.C.; Seralini, G.E. Glyphosate-based herbicides are toxic and endocrine disruptors in human cell lines. *Toxicology* **2009**, *262*, 184–191. [CrossRef] [PubMed]
186. Walsh, L.P.; McCormick, C.; Stocco, D.M. Roundup inhibits steroidogenesis by disrupting steroidogenic acute regulatory (StAR) protein expression. *Environ. Health Perspect.* **2000**, *108*, 769–776. [CrossRef]
187. Thongprakaisang, S.; Thiantanawat, A.; Rangkadilok, N.; Suriyo, T.; Satayavivad, J. Glyphosate induces human breast cancer cells growth via oestrogen receptors. *Food Chem. Toxicol.* **2013**, *59*, 129–136. [CrossRef] [PubMed]
188. Sritana, N.; Suriyo, T.; Kanitwithayanun, J.; Songvasin, B.H.; Thiantanawat, A.; Satayavivad, J. Glyphosate induces growth of oestrogen receptor alpha positive cholangiocarcinoma cells via non-genomic oestrogen receptor/ERK1/2 signaling pathway. *Food Chem. Toxicol.* **2018**, *118*, 595–607. [CrossRef]
189. Stur, E.; Aristizabal-Pachon, A.F.; Peronni, K.C.; Agostini, L.P.; Waigel, S.; Chariker, J.; Miller, D.M.; Thomas, S.D.; Rezzoug, F.; Detogni, R.S.; et al. Glyphosate-based Herbicides at Low Doses Affect Canonical Pathways in Estrogen Positive and Negative Breast Cancer Cell Lines. *PLoS ONE* **2019**, *14*, e0219610. [CrossRef]
190. Armiliato, N.; Ammar, D.; Nezzi, L.; Stralio, M.; Muller, Y.M.R.; Nazari, E.M. Changes in ultrastructure and expression of steroidogenic factor-1 in ovaries of zebrafish *Danio rerio* exposed to glyphosate. *J. Toxicol. Environ. Health A* **2014**, *77*, 405–414. [CrossRef]
191. Xie, L.; Thrippleton, K.; Irwin, M.A.; Siemering, G.S.; Mekebri, A.; Crane, D.; Berry, K.; Schlenk, D. Evaluation of estrogenic activities of aquatic herbicides and surfactants using a rainbow trout vitellogenin assay. *Toxicol. Sci.* **2005**, *87*, 391–398. [CrossRef]
192. Gandhi, J.S.; Cecala, K.K. Interactive effects of temperature and glyphosate on the behavior of Blue Ridge two-lined salamanders (*Eurycea wilderae*). *Environ. Toxicol. Chem.* **2016**, *35*, 2297–2303. [CrossRef]
193. Guilherme, S.; Gaivao, I.; Santos, M.A.; Pacheco, M. DNA and chromosomal damage induced in fish (*Anguilla* L.) by aminomethylphosphonic acid (AMPA) –the major environmental breakdown product of glyphosate. *Environ. Sci. Pollut. Res. Int.* **2014**, *21*, 8730–8739. [CrossRef] [PubMed]
194. Mañas, F.; Peralta, L.; Raviolo, J.; García-Ovando, H.; Weyers, A.; Ugnia, L.; Gonzalez-Cid, M.; Larripa, I.; Gorla, N. Genotoxicity of AMPA, the environmental metabolite of glyphosate, assessed by the Comet assay and cytogenetic tests. *Ecotoxicol. Environ. Saf.* **2009**, *72*, 834–837. [CrossRef] [PubMed]
195. FoEE. Human contamination by glyphosate. Friends of the Earth Europe. 2013. Available online: [https://www.foeeurope.org/sites/default/files/press\\_releases/foee\\_4\\_human\\_contamination\\_glyphosate.pdf](https://www.foeeurope.org/sites/default/files/press_releases/foee_4_human_contamination_glyphosate.pdf) (accessed on 15 June 2020).
196. CFS—Center for Food Safety. *Glyphosate and Cancer Risk: Frequently Asked Questions*; Fact Sheet: Washington, DC, USA, May 2015.
197. Defarge, N.; Takács, E.; Lozano, V.L.; Mesnage, R.; Spiroux de Vendômois, J.; Seralini, G.-E.; Székács, A. Co-formulants in glyphosate-based herbicides disrupt aromatase activity in human cells below toxic levels. *Inter. J. Environ. Res. Public Health* **2016**, *13*, 264. [CrossRef]
198. Acquavella, J.F.; Alexander, B.H.; Mandel, J.S.; Gustin, C.; Baker, B.; Chapman, P.; Bleeke, M. Glyphosate biomonitoring for farmers and their families: Results from the Farm Family Exposure Study. *Environ. Health Perspect.* **2004**, *112*, 321–326. [CrossRef]
199. Mandel, J.S.; Alexander, B.H.; Barker, B.A.; Acquavella, J.F.; Chapman, P.; Honeycutt, R. Biomonitoring for fam families in the farm family exposure study. *Scand. J. Work. Environ. Health* **2005**, *31*, 98–104.
200. Baker, B.A.; Alexander, B.H.; Mandel, J.S.; Acquavella, J.F.; Honeycutt, R.; Chapman, P. Farm Family Exposure Study: Methods and Recruitment Practices for a Biomonitoring Study of Pesticide Exposure. *J. Expo. Anal. Environ. Epidemiol.* **2005**, *15*, 491–499. [CrossRef]
201. McGuire, M.K.; McGuire, M.A.; Price, W.J.; Shafii, B.; Carrothers, J.M.; Lackey, K.A.; Goldstein, D.A.; Jensen, P.K.; Vicini, J.L. Glyphosate and aminomethylphosphonic acid are not detectable in human milk. *Am. J. Clin. Nutr.* **2016**, *103*, 1285–1290. [CrossRef]
202. Niemann, L.; Sieke, C.; Pfeil, R.; Solecki, R. A critical review of glyphosate findings in human urine samples and comparison with the exposure of operators and consumers. *J. Verbr. Lebensm.* **2015**, *10*, 3–12. [CrossRef]
203. Stephenson, C.L.; Harris, C.A. An assessment of dietary exposure to glyphosate using refined deterministic and probabilistic methods. *Food Chem. Toxicol.* **2016**, *95*, 28–41. [CrossRef]

204. Stephenson, C.L.; Harris, C.A.; Clarke, R. An Assessment of the Acute Dietary Exposure to Glyphosate Using Deterministic and Probabilistic Methods. *Food Addit. Contam. A* **2018**, *35*, 258–272. [[CrossRef](#)]
205. EPA. *Available Information on Assessing Exposure from Pesticides in Food: A User's Guide*; EPA-HQ-OPP-2007-0780-0001; Office of Pesticide Programs: Washington, DC, USA, 2000.
206. Curwin, B.D.; Hein, M.J.; Sanderson, W.T.; Striley, C.; Heederik, D.; Kromhout, H.; Reynolds, S.J.; Alavanja, M.C. Urinary pesticide concentrations among children, mothers and fathers living in farm and non-farm households in Iowa. *Ann. Occup. Hyg.* **2007**, *51*, 53–65. [[CrossRef](#)]
207. Curwin, B.D.; Hein, M.J.; Sanderson, W.T.; Striley, C.; Heederik, D.; Kromhout, H.; Reynolds, S.J.; Alavanja, M.C. Pesticide dose estimates for children of Iowa farmers and non-farmers. *Environ. Res.* **2007**, *105*, 307–315. [[CrossRef](#)] [[PubMed](#)]
208. Aris, A.; Leblanc, S. Maternal and fetal exposure to pesticides associated to genetically modified foods in Eastern Townships of Quebec, Canada. *Reprod. Toxicol.* **2011**, *31*, 528–533. [[CrossRef](#)]
209. EPA. *Glyphosate Human Health Risk Assessment for Proposed Use on Indian Mulberry and Amend Use on Pea*; EPA: Washington, DC, USA, 2006; p. 21.
210. Antoniou, M.; Habib, M.; Howard, C.V.; Jennings, R.C.; Leifert, C.; Fagan, J. Teratogenic effects of glyphosate-based herbicides: Divergence of regulatory decisions from scientific evidence. *J. Environ. Anal. Toxicol.* **2012**, *54*, 006. [[CrossRef](#)]
211. Sheppard, L.; Shaffer, R.M. Re: Glyphosate Use and Cancer Incidence in the Agricultural Health Study. *J. Natl. Cancer Inst.* **2019**, *111*, 214–215. [[CrossRef](#)]
212. Connolly, A.; Coggins, M.A.; Koch, H.M. Human biomonitoring of glyphosate exposures: State-of-the-art and future research challenges. *Toxics* **2020**, *8*, 60. [[CrossRef](#)] [[PubMed](#)]
213. National Academies Press. *National Academies of Sciences, Engineering, and Medicine. Genetically Engineered Crops: Experiences and Prospects*; The National Academies Press: Cambridge, MA, USA, 2016. [[CrossRef](#)]
214. Benbrook, C.M. Enhancements needed in GE crop and food regulation in the U.S. *Front. Public Health* **2016**, *4*, 59. [[CrossRef](#)] [[PubMed](#)]
215. Benachour, N.; Seralini, G.E. Glyphosate formulations induce apoptosis and necrosis in human umbilical, embryonic, and placental cells. *Chem. Res. Toxicol.* **2009**, *22*, 97–105. [[CrossRef](#)]
216. Benbrook, C.M. Why regulators lost track and control of pesticide risks: Lessons from the case of glyphosate-based herbicides and genetically engineered-crop technology. *Curr. Environ. Health Rep.* **2018**, *5*, 387–395. [[CrossRef](#)] [[PubMed](#)]
217. Sims, B.; Corsi, S.; Gbehounou, G.; Kienzle, J.; Taguchi, M.; Friedrich, T. Sustainable weed management for conservation agriculture: Options for smallholder farmers. *Agriculture* **2018**, *8*, 118. [[CrossRef](#)]
218. Beckie, H.J.; Flower, K.C.; Ashworth, M.B. Farming without Glyphosate? *Plants* **2020**, *9*, 96. [[CrossRef](#)]
219. FAOSTAT. 2019. Available online: <http://www.fao.org/faostat/en/#data> (accessed on 15 June 2020).
220. Megacity Commission of the International Geographical Union. 2006. Available online: <http://www.megacities.uni-koeln.de/documentation/> (accessed on 15 June 2020).
221. Powles, S.B.; Preston, C. Evolved glyphosate resistance in plants: Biochemical and genetic basis of resistance. *Weed Technol.* **2006**, *20*, 282–289. [[CrossRef](#)]
222. Smith, T.M.; Reynolds, R.W. A global merged land air and surface temperature reconstruction based on historical observations (1880–1997). *J. Clim.* **2005**, *18*, 2021–2036. [[CrossRef](#)]
223. Hansen, S.O.; Hattendorf, J.; Wittenberg, R.; Reznik, S.Y.; Nielsen, C.; Ravn, H.P.; Nentwig, W. Phytophagous insects of giant hogweed *Heracleum mantegazzianum* (Apiaceae) in invaded areas of Europe and in its native area of the Caucasus. *Eur. J. Entomol.* **2006**, *103*, 387–396. [[CrossRef](#)]