



Review

# Humic Acids Aggregates as Microheterogeneous Reaction Media: Alkaline Hydrolysis Reactions

Antonio Cid-Samamed <sup>1,2,3,4,\*</sup> , Luis García-Río <sup>5</sup>, Juan C. Mejuto <sup>4,\*</sup>  and Alejandro Saavedra <sup>4</sup>

<sup>1</sup> i4HB–Associate Laboratory Institute for Health and Bioeconomy, NOVA School of Science and Technology, NOVA University Lisbon, 2819-516 Caparica, Portugal

<sup>2</sup> UCIBIO–Applied Molecular Biosciences Unit, NOVA School of Science and Technology, NOVA University of Lisbon, 2829-516 Caparica, Portugal

<sup>3</sup> LAQV-REQUIMTE, NOVA School of Science and Technology, NOVA University of Lisbon, 2829-516 Caparica, Portugal

<sup>4</sup> Physical Chemistry Department, Faculty of Sciences, University of Vigo, 32004 Ourense, Spain; alexardevaas@gmail.com

<sup>5</sup> Departamento de Química Física, Facultade de Química, Universidade de Santiago de Compostela, 15782 Santiago, Spain; luis.garcia@usc.es

\* Correspondence: acids@fct.unl.pt (A.C.-S.); xmejuto@uvigo.es (J.C.M.); Tel.: +34-988387031 (J.C.M.)

**Abstract:** The influence of humic aggregates in a water solution upon the chemical stability under basic conditions of different substrates was reviewed. The kinetic behavior of each substrate was modeled in terms of a micellar pseudophase model.

**Keywords:** humic acids; alkaline hydrolysis; pseudophase model; micelles



**Citation:** Cid-Samamed, A.; García-Río, L.; Mejuto, J.C.; Saavedra, A. Humic Acids Aggregates as Microheterogeneous Reaction Media: Alkaline Hydrolysis Reactions. *Compounds* **2022**, *2*, 131–143. <https://doi.org/10.3390/compounds2020010>

Academic Editor: Dmitrii S. Bolotin

Received: 27 December 2021

Accepted: 19 April 2022

Published: 28 April 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).


## 1. Introduction

It is well known that colloidal aggregates constitute true chemical nano-reactors in their different forms. By concentrating the reactants in restricted areas or separating them between the different domains of the system, these reaction media can cause both critical catalytic phenomena and dramatic inhibitions of chemical processes [1–3].

One such reaction medium is micellar aggregates. It is well known the influence of aqueous micelles upon the chemical equilibria and the rate of chemical reactions. Evidences of these effects have been summarized in the literature [4–6]. Micellar catalysis on organic and inorganic reactions has been extensively studied, and it is well known that it is related to the ability of micelles to absorb all types of molecules [7–9]. The reagents incorporation into the micelle is driven by hydrophilic/hydrophobic interactions and electrostatic interactions between the micellar surface and the reagents. In this way, the interaction between the reagents and the micellar aggregate can be modeled by applying the pseudophase model [10]. In this model, bulk water can be considered like one continuous phase and micellar aggregate as another. The reagents would be distributed between both phases, or they would be left behind from one of them. In the same way, the chemical reaction could also occur in both phases or in one of them.

We have introduced the kinetic behavior in the presence of micelles because humic acids have behavior in the aqueous solutions similar to micellar aggregates [11,12].

Humic substances (HSs) represent a significant part of the organic matter that is found in soils and in natural environments. These have a high specific surface area [13–15], with a negative surface charge due to numerous deprotonable carboxylic and phenolic groups (see Scheme 1) [16]. Furthermore, due to amino groups that can be protonated, it enables a positive surface charge. That is why the surface charge has a significant dependence on pH. Full ionized charge capacity (CEC) is 0.3–0.6 mol·kg<sup>−1</sup>.



**Scheme 1.** Model of the humic acid chemical structure.

The basic hydrolysis reaction kinetics of the reagents assessed in this review paper were followed by the first-order conditions ( $[V] \ll [\text{OH}^-]$ ), using for this aim Varian Cary 50 Bio spectrophotometer with the thermostated reader cell at  $(25.0 \pm 0.1)^\circ\text{C}$ . The HSs used for these experiments absorb in the UV-vis region, so the spectra were carried out from 200 to 800 nm, using as blank the spectrum of HSs without reaction (in absence of the reagent to study). As previously reported, the disappearance of the shoulder at maximum wavelength was critical to assess the basic hydrolysis of the molecules tested (e.g.,  $\lambda_{\text{max(VI)}} = 242 \text{ nm}$ ,  $\lambda_{\text{max(HCF)}} = 275 \text{ nm}$ ). The data were recorded in triplicate to ensure the reproducibility of the method. As abovementioned, the data were treated by assuming the conditions of the pseudo-first order and the rate equation to determine the reagents R disappearance as follows:

$$-\frac{d[R]}{dt} = k_w[\text{Rs}]_t[\text{OH}^-]_t = k_{\text{obs}}[\text{Rs}]_t \quad (1)$$

where  $[R]$  is the reagents concentration,  $k_w$  is the bimolecular rate, and  $k_{\text{obs}}$  is the pseudo-first rate constants for the basic hydrolysis of the reagents.

In aqueous solution [17–21], in which HSs acts as micellar-like aggregates, they can adsorb different substrates [22–29] that can act as environmental pollutants [30], increasing or reducing their bioavailability, which is the case of different xenobiotics [31,32] that are persistent in the aquatic environment or soils [33,34]. The role of humates and humic acids in different organic substrates' mobility was conveniently illustrated by Ramus et al. [35,36] and Chiou et al. [37]. Thus, it was shown that when the content of humic substances in solution increases, significant decreases are observed in the mass transfer values at the gas–liquid interface. Furthermore, they can modify pollutants' destination in the environment and act as catalysts in the chemical breakdown of some pollutants [31,38–44]. Previous studies show the effect of HSs in solution on hydrolysis reactions (the main transformation pathway for many xenobiotics in the environment [39,45]). These substrates are typically composed of a marked hydrophobic character that shows a high affinity for HSs aggregates [38–46].


Humic substances are also able to “kidnap” heavy metals [47–54] and interact with minerals [28,49,55,56], modifying the absorption and toxicity of these compounds. There are few studies in the literature in which the catalytic activity of HSs has been compared [25,38,45,46,57,58]. These studies can be critical, considering their possibility in the biogeochemical cycle of elements and the high concentrations of HSs in some natural environments.

In this paper, we will review some of the contributions of our research group [59–63], which has tried to shed light on the role of humic acids and humates in aqueous solution,

as colloidal aggregates, in the catalysis of different basic hydrolysis processes. HSs used in these studies were isolated from soil using the method previously described [64].

## 2. Hydrolysis of Carbocations

The first group of reactions that we decided to analyze in humic substances was the alkaline fading of stable triaryl methyl carbocations (Scheme 2) [61]. This is because it is a reaction that has already been studied, and these carbocations entangled with different nucleophiles were used to construct of the nucleophilic scale of the Ritchie N<sup>+</sup> family [65,66], which was considered as a challenge to the reactivity–selectivity principle [67,68]. These reasons made it an instrument for studying reactivity in other microorganized environments. One of the first published studies on catalysis was carried out precisely with the basic hydrolysis of crystal violet [69,70]. Subsequently, they were widely used in studies on the catalytic or inhibitory effects of normal micelles [71], reverse micelles or microemulsions [72–74], and other microheterogeneous systems [75]. In this way, we understood that it could be an excellent chemical probe to obtain valuable information on the role of the different factors that affect the general reactivity in HSs micelles, such as the reagents' compartmentalization and the characteristics of the HSs aggregates as reaction medium.




**Scheme 2.** Crystal violet (CV) and malachite green (MG).

Thereby, the basic hydrolysis of CV and MG in the presence of different concentrations of HSs has been studied under conditions in which all the humic acids are in the form of humate. The presence of HSs has been observed to inhibit the hydrolysis reaction, as shown in Table 1. An inhibition of 4.5 times was observed for the hydrolysis of CV and 24 times for the case of MG.

**Table 1.** Kinetic parameters of basic hydrolysis of carbocations in the presence of HSs aggregates obtained from the pseudophase model (Equation (2)).

Substrate	$k_w/M^{-1} \cdot \text{min}^{-1}$	$K_s/\text{mg}^{-1} \cdot \text{L}$
CV [61]	$1.23 \pm 0.02$	$0.13 \pm 0.01$
MG [61]	$181 \pm 6$	$0.65 \pm 0.05$

This behavior is justified based on the distribution of the reagents and the different reaction loci in the microheterogeneous system. Thus, the reaction can occur inside the aggregate, on the aggregate's surface (in the Stern layer), or in the bulk solvent. The interior of the aggregate will have a hydrophobic nature. However, the CV or MG housed there could not be exposed to OH<sup>−</sup> and therefore would not participate in the reaction. The same would happen on the aggregate's surface since the OH<sup>−</sup> would also be excluded due to electrostatic repulsions between the OH<sup>−</sup> and the negatively charged groups located on the surface of the aggregate [76,77]. Thus, they only take place in the bulk water, where the OH<sup>−</sup> and a part of the carbocations non-associated with the aggregate will be found. Scheme 3 shows the mechanism that takes place in this microheterogeneous system.



**Scheme 3.** Pseudophase model for basic hydrolysis of CV/MG in the presence of HSs.


The observed results [61] would be equivalent to those observed in anionic micelles [78]. The observed kinetic behavior complies with the kinetic equation obtained from Scheme 3 (Equation (2)), where the subscripts *s* and *w* denote the pseudophase micellar and the bulk water, respectively, and  $K_s$  (Equation (3)) is the carbocation association constant to the micellar aggregate, and  $k_2$  is the rate constant in pure water.

$$k_2 = \frac{k_w}{1 + K_s[\text{HSs}]} \quad (2)$$

$$K_s = \frac{[R]_s}{[R]_w[\text{HSs}]} \quad (3)$$

$$\frac{1}{k_2} = \frac{1}{k_w} + \frac{K_s}{K_w}[\text{HSs}] \quad (4)$$


From these equations, we obtained values of the substrate association constants to HSs aggregate. The values obtained were  $K_s = (0.13 \pm 0.01) \text{ mg}^{-1} \cdot \text{L}$  and  $K_s = (0.65 \pm 0.05) \text{ mg}^{-1} \cdot \text{L}$ , respectively, for the CV and the MG. This higher value of  $K_s$  observed for the MG versus the CV is due to the lower polarity of the first versus the second, with which it would be expected that it would penetrate more deeply into the HSs [60]. Figure 1 shows the fit of experimental results to linearized Equation (2) (Equation (4)). Table 1 shows the kinetic parameter for these reactions.



**Figure 1.** Pseudophase model (Equation (4)) fit of experimental results for basic hydrolysis of CV (○) and MG (●) in the presence of humic aggregates.  $T = 25 \text{ }^\circ\text{C}$ .


### 3. Hydrolysis of N-Methyl-N-nitroso-p-toluene Sulfonamide

Another molecule of interest whose study was approached in our laboratory was N-methyl-N-nitroso-p-toluenesulfonamide (MNTS) [54] (see Scheme 4), which is of biological interest [79] because it is well known that nitrosulfonamides are very effective nitrosating agents [68]. Another advantage of this molecule is the detailed research of our team of the mechanisms of its hydrolysis reactions (both acidic and basic) together with the mechanisms of the transnitrosation processes that involve it, both in homogeneous and microheterogeneous media [1–3,5,80–83]. That is why we consider it as a suitable chemical probe for its study in aggregated HSs since it could be a valuable tool for deepening the knowledge of the chemical reactivity in the presence of HSs micelles, complementing the previous observations acquired from the study of the processes of hydrolysis of CV and MG (*vide supra*).



**Scheme 4.** N-methyl-N-nitroso-p-toluenesulfonamide (MNTS).


As in the previous section, the presence of aggregated HSs inhibits the basic hydrolysis of MNTS by the association of MNTS with the aggregates. Again, this behavior was modeled using the pseudophase model. Considerations like those applied for CV, and MG, led us to propose a mechanism like that of the previous ionic compounds, such as the one shown in Scheme 3, so that, once again, the corresponding kinetic equation would be Equation (2) and the distribution of the MNTS would be governed by Equation (3). From the adjustment of the experimental results to Equation (2), a value for  $K_s = (25 \pm 2) \text{ mg}^{-1} \cdot \text{L}$  was obtained (see Figure 2).



**Figure 2.** Pseudophase model (Equation (4)) fit of experimental results for basic hydrolysis of MNTS in the presence of humic aggregates.  $T = 25 \text{ }^\circ\text{C}$ .

#### 4. Hydrolysis of Carbofuran and Derivatives, Iprodione, and Vinclozolin

Previous results show that the comparability presented by the HSs aggregates have a kinetic behavior analogous to the conventional micellar aggregates. With the study of the basic hydrolysis of both ionic substances (MG and CV) [60] and with non-ionic substances (MNTS [54]), the validity of the micellar pseudophase model was demonstrated beyond any doubt. For this reason, the stability of different xenobiotics in basic media and in the presence of HSs in aqueous solution was discussed below. In this way, the basic hydrolysis of carbofuran (CF) [62] and two derivatives of carbofuran [62]—3-hydroxy-carbofuran (HCF) and 3-keto-carbofuran (KCF), as well as iprodione (IP) [61] and vinclozolin (VI) [59] in the presence of HSs, were analyzed, and their behavior was compared with similar results in ionic and non-ionic micelles. Scheme 5 shows the xenobiotics under study.




**Scheme 5.** Carbofuran and carbofuran derivatives, iprodione (IP), and vinclozolin (VI).

In the case of carbamates, a curious behavior was observed since we found an inhibition on the basic hydrolysis of HCF [62] and KCF [62], justifiable, as in the previous cases, based on the association of the substrates to the aggregate and the exclusion of  $\text{OH}^-$  from their vicinity based on electrostatic considerations (*vide supra*). When the experimental data is fitted to a mechanism similar to that used for the case of MNTS and ionic compounds, (Scheme 3) (Equations (2) and (3)) we obtain values of the constant in water of  $k_w = (1.86 \pm 0.06) \times 10^2 \text{ M}^{-1} \cdot \text{min}^{-1}$  and  $k_w = (11.4 \pm 0.6) \times 10^3 \text{ M}^{-1} \cdot \text{min}^{-1}$  and some values of association constants of  $K_s = (1.0 \pm 0.1) \times 10^{-2} \text{ mg}^{-1} \cdot \text{L}$  and  $K_s = (5 \pm 1) \times 10^{-3} \text{ mg}^{-1} \cdot \text{L}$ , respectively, for HCF and KCF. However, no effect is found on the reaction rate for CF [62] hydrolysis, so we must assume that in this case, CF absorption does not occur inside the HSs aggregate, and all the CF remains in the bulk water. Table 2 shows the kinetic results obtained and Figure 3, as an example, shows the pseudophase model fit of experimental results for basic hydrolysis of carbofuran and carbofuran-derivatives.

**Table 2.** Kinetic parameters of basic hydrolysis different xenobiotics in the presence of HSs aggregates obtained from the pseudophase model (Equation (2)).

Substrate	$k_w/M^{-1}\cdot\text{min}^{-1}$	$K_s/\text{mg}^{-1}\cdot\text{L}$
CF [62]	$(0.66 \pm 0.06) \times 10^2$	0
HCF [62]	$(1.86 \pm 0.06) \times 10^2$	$(1.0 \pm 0.1) \times 10^{-2}$
KCF [62]	$(11.40 \pm 0.60) \times 10^3$	$(5.0 \pm 1.0) \times 10^{-3}$
IP [61]	$(1.88 \pm 0.01) \times 10^3$	$(1.4 \pm 0.1) \times 10^{-2}$
VI [59]	$(8.20 \pm 0.40) \times 10^2$	$(9.7 \pm 0.1) \times 10^{-2}$

**Figure 3.** Pseudophase model (Equation (4)) fit of experimental results for basic hydrolysis of CF (○), HCF (●), and KCF (□) in the presence of humic aggregates.  $T = 25\text{ }^{\circ}\text{C}$ .

When the basic hydrolysis of iprodione (IP) [61] was analyzed, inhibition was also observed. This behavior indicates that the mechanistic model applied in the case of carbofuran and carbofuran-derivatives is still valid. In this case, a slightly higher inhibition was observed than in the previous cases. In the case of carbofuran-derivatives, an inhibition of ~1.7- and 1.5-fold, respectively, had been found for the case of HCF and KCF (we must remember that we had not found an effect of the presence of HSs aggregates on the basic hydrolysis of CF) while in the case of the IP, the decrease in speed was 2-fold.

As in the previous cases, from Equation (2), we have obtained the corresponding kinetic parameters. A rate constant for the hydrolysis process in bulk water of  $k_w = (1.878 \pm 0.006) \times 10^3\text{ M}^{-1}\cdot\text{min}^{-1}$  and an association constant to the addition of  $K_s = (1.40 \pm 0.10) \times 10^{-2}\text{ mg}^{-1}\cdot\text{L}$  (Table 2). The higher value of the observed  $K_s$  would justify the greater inhibition observed on the hydrolysis of the IP compared to the carbofuran-derivatives.

Regarding the results obtained for the basic hydrolysis of vinclozolin (VI) [59], the inhibition observed due to the presence of HSs aggregates was much more dramatic, obtaining a decrease in the reaction rate of 9-fold. When the basic hydrolysis of iprodione (IP) was analyzed, inhibition was also observed. By applying the pseudophase model and adjusting the experimental data to Equation (2), we obtain an association constant to the aggregate significantly higher than in the previous cases,  $K_s = (9.7 \pm 0.1) \times 10^{-2}\text{ mg}^{-1}\cdot\text{L}$  (Table 2). The greater affinity of the VI towards the HSs aggregates would cause this more significant inhibition observed on the basic hydrolysis in this compound.




## 5. Binding Constants and Hydrophobicity of HSs Core

Because of the units of  $K_s$ , it is not possible to apply a direct comparison of this value with the corresponding ones for normal micelles to evaluate the HSs core's hydrophobicity.

However, with the data obtained for the different hydrolysis reactions discussed in the previous sections, we can compare the ratios between the association constants obtained for aggregated HSs and normal micelles.


In the literature, there are abundant studies that obtain the association constants of some of these compounds (MNTS [82,84–86] and CV [75,87]) to a large number of micelle aggregates, finding a linear correlation between  $K_s(\text{MNTS})/K_s(\text{CV})$  and the length of the hydrocarbon chain of traditional surfactants. Thus, the  $K_s(\text{MNTS})/K_s(\text{CV})$  ratio in the presence of HSs aggregates is 192. For the OTACl (octadecyl trimethyl ammonium chloride), we find that  $K_s(\text{MNTS})/K_s(\text{CV})$  has a value of 10, that is, 19 times less than that found for HSs. In the case of SDS (sodium dodecyl sulfate), the value of  $K_s(\text{MNTS})/K_s(\text{CV})$  would have a value of 4.3, in this case, 45 times less than the corresponding one for HSs. Figure 4 shows the ratio value between MNTS and CV binding constants to different chain length surfactants. It is, therefore, evident that the core of aggregate HSs is drastically more hydrophobic than that of traditional micelles, both anionic and cationic.



**Figure 4.** Binding constants ratio between MNTS and CV to normal micelles as a function of the chain length of surfactants.  $T = 25\text{ }^\circ\text{C}$ .

Furthermore, it is evident that the hydrophilic/hydrophobic interactions are the main driving force behind the association of the different substrates to the aggregate, as evidenced by the evident correlation between the  $\log P$  values of the substrates and the association constants found from the adjustment of the experimental data to the micellar pseudophase model (see Figure 5).





**Figure 5.** Relationship between  $\log K_s$  for different substrates (KCF, HCF, IP, MNTS, VI) in humic aggregates and the  $\log P$  value for each substrate.  $T = 25\text{ }^\circ\text{C}$ .

## 6. Conclusions

In summary, the effect of the presence of micelles formed by HSs on the basic hydrolysis reactions results in an inhibition due to the exclusion of  $\text{OH}^-$  from the Stern layer of the aggregate due to the electrostatic interactions between the negative surface charge of the HSs and the  $\text{OH}^-$  ion. In this way, the inhibitions would be due to the inclusion of the different substrates inside micelles, protecting them from a nucleophilic attack by  $\text{OH}^-$ .

The hydrophilic/hydrophobic interactions are the main driving force behind the association of the different substrates to the aggregate, which has been demonstrated due to the evident correlation between the  $\log P$  values of the different substrates and their binding constants.

Therefore, when comparing the constants obtained in the presence of HSs with the constants of association to other micellar aggregates, it can be concluded that the core of the aggregates of HSs has a hydrophobic character significantly greater than that of “usual” micelles of sodium alkyl sulfate, alkyl trimethyl ammonium chloride or alkyl pyridinium chloride, even for hydrocarbon chains of more than 18 carbon atoms.

**Author Contributions:** Conceptualization, J.C.M. and L.G.-R.; resources, A.C.-S. and A.S.; writing—original draft preparation, J.C.M. and A.C.-S.; writing—review and editing, L.G.-R. and J.C.M.; visualization, A.C.-S. and A.S.; supervision, J.C.M.; project administration, J.C.M.; funding acquisition, L.G.-R. and J.C.M. All authors have read and agreed to the published version of the manuscript.

**Funding:** Financial support from the Ministerio de Economía y Competitividad of Spain (project CTQ2017-84354-P), Xunta de Galicia and the European Regional Development Fund (ERDF) is gratefully acknowledged (GR 2007/085 and ED431C2018/42-GRC). A.C.-S. thanks to FCT-Fundação para a Ciência e a Tecnologia, I.P., in the scope of the project UIDP/04378/2020 and UIDB/04378/2020 of the Research Unit on Applied Molecular Biosciences-UCIBIO and the project LA/P/0140/2020 of the Associate Laboratory Institute for Health and Bioeconomy-i4HB.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Conflicts of Interest:** The authors declare no conflict of interest, and the funders had no role in the design of the study, in the collection, analyses, or interpretation of data, in the writing of the manuscript, or in the decision to publish the results.

## Abbreviations

CEC	ionized charge capacity
CF	carbofuran
CV	crystal violet
HCF	3-hydroxy-carbofuran
HSs	humic substances
IP	iprodione
KCF	3-keto-carbofuran
MG	malachite green
MNTS	N-methyl-N-nitroso-p-toluenesulfonamide
OTACl	octadecyl trimethyl ammonium chloride
SDS	sodium dodecyl sulfate
VI	vinclozolin

## References

1. García-Río, L.; Hervés, P.; Mejuto, J.C.; Pérez-Juste, J.; Rodríguez-Dafonte, P. Comparative study of nitroso group transfer in colloidal aggregates: Micelles, vesicles and microemulsions. *New J. Chem.* **2003**, *27*, 372–380. [[CrossRef](#)]
2. García-Río, L.; Hervés, P.; Mejuto, J.C.; Pérez-Juste, J.; Rodríguez-Dafonte, P. Pseudophase Approach to Reactivity in Microemulsions: Quantitative Explanation of the Kinetics of the Nitroso Group Transfer Reactions between N-methyl-N-nitroso-p-toluenesulfonamide and Secondary Alkylamines in Water/AOT/Isooctane Microemulsions. *Ind. Eng. Chem. Res.* **2003**, *42*, 5450–5456. [[CrossRef](#)]
3. García-Río, L.; Hervés, P.; Mejuto, J.C.; Rodríguez-Dafonte, P. Nitrosation reactions in water/AOT/xylene microemulsions. *Ind. Eng. Chem. Res.* **2006**, *45*, 2. [[CrossRef](#)]
4. Bunton, C.A.; Savelli, G. Organic Reactivity in Aqueous Micelles and Similar Assemblies. *Adv. Phys. Org. Chem.* **1986**, *22*, 213–309.
5. García-Río, L.; Hervés, P.; Leis, J.R.; Mejuto, J.C.; Rodríguez-Dafonte, P. Reactive micelles: Nitroso group transfer from N-methyl-N-nitroso-p-toluenesulfonamide to amphiphilic amines. *J. Phys. Org. Chem.* **2004**, *17*, 1067–1072. [[CrossRef](#)]
6. Huang, X.; Xue, L. Recent Progress in Enzyme Catalysis in Reverse Micelles. In *Bridging Heterogeneous and Homogeneous Catalysis: Concepts, Strategies, and Applications*; Wiley-VCH: Weinheim, Germany, 2014; ISBN 9783527675906.
7. Bunton, C.A. Micellar rate effects-assumptions and approximations. *Arkivoc* **2011**, *2011*, 490–504. [[CrossRef](#)]
8. Bunton, C.A.; Yatsimirsky, A.K. Comparisons and Analyses of Theoretical Treatments of Micellar Effects upon Ion–Molecule Reactions. Relevance to Amide Exchange. *Langmuir* **2000**, *16*, 5921–5931. [[CrossRef](#)]
9. Bunton, C.A. The dependence of micellar rate effects upon reaction mechanism. *Adv. Colloid Interface Sci.* **2006**, *123–126*, 333–343. [[CrossRef](#)]
10. Bunton, C.A. Reactivity in aqueous association colloids. Descriptive utility of the pseudophase model. *J. Mol. Liq.* **1997**, *72*, 231–249. [[CrossRef](#)]
11. Tian, S.; Tan, W.; Wang, X.; Li, T.; Song, F.; Huang, N.; Bai, Y. Surface Activity of Humic Acid and Its Sub-Fractions from Forest Soil. *Sustainability* **2021**, *13*, 8122. [[CrossRef](#)]
12. Arias, M.; García-Río, L.; Mejuto, J.C.; Rodríguez-Dafonte, P.; Simal-Gándara, J. Influence of Micelles on the Basic Degradation of Carbofuran. *J. Agric. Food Chem.* **2005**, *53*, 7172–7178. [[CrossRef](#)] [[PubMed](#)]
13. Flis-Bujak, M.; Ksiezopolska, A.; Stawinski, J.; Turski, R. The surface properties of humic acids extracted from chernozem and grey-brown podzolic soils. *Int. Agrophysics* **1995**, *9*, 115–123.
14. Mendez, J.C.; Hiemstra, T.; Koopmans, G.F. Assessing the Reactive Surface Area of Soils and the Association of Soil Organic Carbon with Natural Oxide Nanoparticles Using Ferrihydrite as Proxy. *Environ. Sci. Technol.* **2020**, *54*, 11990–12000. [[CrossRef](#)] [[PubMed](#)]
15. Chiou, C.T.; Lee, J.F.; Boyd, S.A. The surface area of soil organic matter. *Environ. Sci. Technol.* **1990**, *24*, 1164–1166. [[CrossRef](#)]
16. De Melo, B.A.G.; Motta, F.L.; Santana, M.H.A. Humic acids: Structural properties and multiple functionalities for novel technological developments. *Mater. Sci. Eng. C* **2016**, *62*, 967–974. [[CrossRef](#)]
17. Conte, P.; Piccolo, A. Conformational arrangement of dissolved humic substances. Influence of solution composition on association of humic molecules. *Environ. Sci. Technol.* **1999**, *33*, 1682–1690. [[CrossRef](#)]
18. Clapp, C.E.; Hayes, M.H.B. Sizes and shapes of humic substances. *Soil Sci.* **1999**, *164*, 777–789. [[CrossRef](#)]
19. He, W.; Bai, Z.; Li, Y.; Kong, X.; Liu, W.; Yang, C.; Yang, B.; Xu, F. Advances in environmental behaviors and effects of dissolved organic matter in aquatic ecosystems. *Sci. China Earth Sci.* **2016**, *59*, 746–759. [[CrossRef](#)]
20. Arias-Estévez, M.; López-Periago, E.; Martínez-Carballo, E.; Simal-Gándara, J.; Mejuto, J.C.; García-Río, L. The mobility and degradation of pesticides in soils and the pollution of groundwater resources. *Agric. Ecosyst. Environ.* **2008**, *123*, 247–260. [[CrossRef](#)]
21. Rutherford, D.W.; Chiou, C.T.; Kille, D.E. Influence of soil organic matter composition on the partition of organic compounds. *Environ. Sci. Technol.* **1992**, *26*, 336–340. [[CrossRef](#)]

22. Leboeuf, E.J.; Weber, W.J., Jr. Macromolecular characteristics of natural organic matter. 2. Sorption and desorption behavior. *Environ. Sci. Technol.* **2000**, *34*, 3632–3640. [[CrossRef](#)]
23. Arias-Estévez, M.; Fernandez-Gandara, D.; Garcia-Falcon, M.S.; Garcia-Rio, L.; Mejuto, J.C.; Simal-Gandara, J. Sorption of PAHs to colloid dispersions of humic substances in water. *Bull. Environ. Contam. Toxicol.* **2007**, *79*, 251–254. [[CrossRef](#)] [[PubMed](#)]
24. Prosen, H.; Zupančič-Kralj, L. The interaction of triazine herbicides with humic acids. *Chromatographia* **2000**, *51*, S155–S164. [[CrossRef](#)]
25. Martin-Neto, L.; Traghetta, D.G.; Vaz, C.M.P.; Crestana, S.; Sposito, G. On the interaction mechanisms of atrazine and hydroxyatrazine with humic substances. *J. Environ. Qual.* **2001**, *30*, 520–525. [[CrossRef](#)]
26. Loiseau, L.; Barriuso, E. Characterization of the atrazine's bound (nonextractable) residues using fractionation techniques for soil organic matter. *Environ. Sci. Technol.* **2002**, *36*, 683–689. [[CrossRef](#)] [[PubMed](#)]
27. Kopinke, F.-D.; Ramus, K.; Poerschmann, J.; Georgi, A. Kinetics of desorption of organic compounds from dissolved organic matter. *Environ. Sci. Technol.* **2011**, *45*, 10013–10019. [[CrossRef](#)] [[PubMed](#)]
28. Arias, M.; Barral, M.T.; da Silva-Carvalho, J.; Mejuto, J.C.; Rubinos, D. Interaction of Hg(II) with kaolin-humic acid complexes. *Clay Miner.* **2004**, *39*, 35–45. [[CrossRef](#)]
29. Chiou, C.T.; McGroddy, S.E.; Kile, D.E. Partition Characteristics of Polycyclic Aromatic Hydrocarbons on Soils and Sediments. *Environ. Sci. Technol.* **1998**, *32*, 264–269. [[CrossRef](#)]
30. Kile, D.E.; Chiou, C.T. Water solubility enhancements of DDT and trichlorobenzene by some surfactants below and above the critical micelle concentration. *Environ. Sci. Technol.* **1989**, *23*, 832–838. [[CrossRef](#)]
31. Prosen, H.; Zupančič-Kralj, L. Evaluation of photolysis and hydrolysis of atrazine and its first degradation products in the presence of humic acids. *Environ. Pollut.* **2005**, *133*, 517–529. [[CrossRef](#)]
32. Noblet, J.A.; Smith, L.A.; Suffet, I.H. Influence of Natural Dissolved Organic Matter, Temperature, and Mixing on the Abiotic Hydrolysis of Triazine and Organophosphate Pesticides. *J. Agric. Food Chem.* **1996**, *44*, 3685–3693. [[CrossRef](#)]
33. Maurya, P.K.; Malik, D.S. Bioaccumulation of xenobiotics compound of pesticides in riverine system and its control technique: A critical review. *J. Ind. Pollut. Control* **2016**, *32*, 580–594.
34. Vigneshwaran, S.; Sirajudheen, P.; Karthikeyan, P.; Nabeena, C.P.; Meenakshi, S. Remediation of Persistent Organic Pesticides from Wastewater Matrices—Present and Future Conceptions. In *Energy, Environment, and Sustainability*; Springer: Singapore, 2021. [[CrossRef](#)]
35. Ramus, K.; Kopinke, F.-D.; Georgi, A. Influence of dissolved humic substances on the mass transfer of organic compounds across the air-water interface. *Chemosphere* **2012**, *86*, 138–143. [[CrossRef](#)]
36. Ramus, K.; Kopinke, F.-D.; Georgi, A. Sorption-induced effects of humic substances on mass transfer of organic pollutants through aqueous diffusion boundary layers: The example of water/air exchange. *Environ. Sci. Technol.* **2012**, *46*, 2196–2203. [[CrossRef](#)] [[PubMed](#)]
37. Chiou, C.T.; Kile, D.E.; Brinton, T.I.; Malcolm, R.L.; Leenheer, J.A.; MacCarthy, P. A comparison of water solubility enhancements of organic solutes by aquatic humic materials and commercial humic acids. MacCarthy. *Environ. Sci. Technol.* **1987**, *21*, 1231–1234. [[CrossRef](#)]
38. Georgi, A.; Reichl, A.; Trommler, U.; Kopinke, F.-D. Influence of sorption to dissolved humic substances on transformation reactions of hydrophobic organic compounds in water. I. Chlorination of PAHs. *Environ. Sci. Technol.* **2007**, *41*, 7003–7009. [[CrossRef](#)]
39. Georgi, A.; Trommler, U.; Reichl, A.; Kopinke, F.-D. Influence of sorption to dissolved humic substances on transformation reactions of hydrophobic organic compounds in water. Part II: Hydrolysis reactions. *Chemosphere* **2008**, *71*, 1452–1460. [[CrossRef](#)]
40. Chiou, C.T.; Malcolm, R.L.; Brinton, T.I.; Kile, D.E. Water solubility enhancement of some organic pollutants and pesticides by dissolved humic and fulvic acids. *Environ. Sci. Technol.* **1986**, *20*, 502–508. [[CrossRef](#)]
41. Chiou, C.T. *Roles of Organic Matter, Minerals, and Moisture in Sorption of Nonionic Compounds and Pesticides by Soil*; Humic Substances in Soil and Crop Sciences: Selected Readings; American Society of Agronomy and Soil Science Society of America: Madison, WI, USA, 2015; pp. 111–160.
42. Chiou, C.T.; Shoup, T.D.; Porter, P.E. Mechanistic roles of soil humus and minerals in the sorption of nonionic organic compounds from aqueous and organic solutions. *Org. Geochem.* **1985**, *8*, 9–14. [[CrossRef](#)]
43. Hung, W.-N.; Lin, T.-F.; Chiu, C.-H.; Chiou, C.T. On the use of a freeze-dried versus an air-dried soil humic acid as a surrogate of soil organic matter for contaminant sorption. *Environ. Pollut.* **2012**, *160*, 125–129. [[CrossRef](#)]
44. Chiou, C.T.; Kile, D.E.; Rutherford, D.W.; Sheng, G.; Boyd, S.A. Sorption of Selected Organic Compounds from Water to a Peat Soil and Its Humic-Acid and Humin Fractions: Potential Sources of the Sorption nonlinearity. *Environ. Sci. Technol.* **2000**, *34*, 1254–1258. [[CrossRef](#)]
45. Singh, S.; Kumar, V.; Singh, J. Kinetic study of the biodegradation of glyphosate by soil bacterial isolates in presence of humic acid, Fe(III) and Cu(II) ions. *J. Environ. Chem. Eng.* **2019**, *7*, 103098. [[CrossRef](#)]
46. Salvestrini, S.; Capasso, S.; Iovino, P. Catalytic effect of dissolved humic acids on the chemical degradation of phenylurea herbicides. *Pest. Manag. Sci.* **2008**, *64*, 768–777. [[CrossRef](#)] [[PubMed](#)]
47. Georgi, A.; Kopinke, F.-D. Validation of a modified flory-huggins concept for description of hydrophobic organic compound sorption on dissolved humic substances. *Environ. Toxicol. Chem.* **2002**, *21*, 1766–1774.

48. Xie, Y.; Guan, Z.; Cheng, J.; Zhou, Y.; Yan, M. Competition between Al(III) and Fe(III) for binding onto natural organic matter: In situ monitoring by UV–Vis absorbance spectroscopy. *Chemosphere* **2021**, *270*, 128655. [[CrossRef](#)]
49. Qu, C.; Chen, W.; Hu, X.; Cai, P.; Chen, C.; Yu, X.-Y.; Huang, Q. Heavy metal behaviour at mineral-organo interfaces: Mechanisms, modelling and influence factors. *Environ. Int.* **2019**, *131*, 104995. [[CrossRef](#)]
50. Marsac, R.; Banik, N.; Lützenkirchen, J.; Catrouillet, C.; Marquardt, C.; Johannesso, K. Modeling metal ion-humic substances complexation in highly saline conditions. *Appl. Geochem.* **2017**, *79*, 52–64. [[CrossRef](#)]
51. Marsac, R.; Davranche, M.; Gruau, G.; Dia, A.; Bouhnik-Le Coz, M. Aluminium competitive effect on rare earth elements binding to humic acid. *Geochim. Et Cosmochim. Acta* **2012**, *89*, 1–9. [[CrossRef](#)]
52. Tipping, E.; Rey-Castro, C.; Bryan, S.E.; Hamilton-Taylor, J. Al(III) and Fe(III) binding by humic substances in freshwaters, and implications for trace metal speciation. *Geochim. Cosmochim. Acta* **2002**, *66*, 3211–3224. [[CrossRef](#)]
53. Esfandiari, N.; Suri, R.; McKenzie, E.R. Competitive sorption of Cd, Cr, Cu, Ni, Pb and Zn from stormwater runoff by five low-cost sorbents; Effects of co-contaminants, humic acid, salinity and pH. *J. Hazard. Mater.* **2022**, *423*, 126938. [[CrossRef](#)]
54. Zhou, P.; Yan, H.; Gu, B. Competitive complexation of metal ions with humic substances. *Chemosphere* **2005**, *58*, 1327–1337. [[CrossRef](#)] [[PubMed](#)]
55. Wang, Y.; Michel, F.M.; Choi, Y.; Eng, P.J.; Levard, C.; Siebner, H.; Gu, B.; Bargar, J.R.; Brown, G.E., Jr. Pb, Cu, and Zn distributions at humic acid-coated metal-oxide surfaces. *Geochim. Cosmochim. Acta* **2016**, *188*, 407–423. [[CrossRef](#)]
56. Arias, M.; Barral, M.T.; Mejuto, J.C. Enhancement of copper and cadmium adsorption on kaolin by the presence of humic acids. *Chemosphere* **2002**, *48*, 1081–1088. [[CrossRef](#)]
57. Klavins, M.; Babre, K. Decarboxylation and alkaline colour fading reactions in presence of humic substances. *Chemosphere* **2002**, *49*, 685–689. [[CrossRef](#)]
58. Kaviš, M.; Dipne, J.; Babre, K. Humic substances as catalysts in condensation reactions. *Chemosphere* **2001**, *44*, 737–742.
59. Morales, J.; Cid, A.; Mejuto, J.C. Alkaline hydrolysis of vinclozolin: Effect of humic acids aggregates in water. *J. Mol. Catal. A Chem.* **2015**, *401*, 13–17. [[CrossRef](#)]
60. Arias-Estevez, M.; Astray, G.; Cid, A.; Fernández-Gándara, D.; García-Río, L.; Mejuto, J.C. Influence of colloid suspensions of humic acids upon the alkaline fading of carbocations. *J. Phys. Org. Chem.* **2008**, *21*, 555–560. [[CrossRef](#)]
61. Morales, J.; Manso, J.A.; Cid, A.; Mejuto, J.C. Stability study of Iprodione in alkaline media in the presence of humic acids. *Chemosphere* **2013**, *92*, 1536–1541. [[CrossRef](#)]
62. Morales, J.; Manso, J.A.; Cid, A.; Mejuto, J.C. Degradation of carbofuran and carbofuran-derivatives in presence of humic substances under basic conditions. *Chemosphere* **2012**, *89*, 1267–1271. [[CrossRef](#)]
63. Astray, G.; García-Río, L.; Lodeiro, C.; Mejuto, J.C.; Moldes, O.; Morales, J.; Moyano, F. Influence of colloid suspensions of humic acids on the alkaline hydrolysis of N-methyl-N-nitroso-p-toluene sulfonamide. *Int. J. Chem. Kinet.* **2010**, *42*, 316–322. [[CrossRef](#)]
64. Klute, A.; Page, A.L. *Methods of Soil Analysis*; American Society of Agronomy: Madison, WI, USA, 1986.
65. Ritchie, C.D. Nucleophilic Reactivities toward Cations. *Acc. Chem. Res.* **1972**, *5*, 348–354. [[CrossRef](#)]
66. Ritchie, C.D.; Virtanen, P.O.I. Cation–Anion Combination Reactions. V. Correlation of Reactivities of Nucleophiles with Diazonium Ions and with Triarylmethyl Cations. *J. Am. Chem. Soc.* **1972**, *94*, 1589–1594. [[CrossRef](#)]
67. Mayr, H.; Ofial, A.R. The reactivity-selectivity principle: An imperishable myth in organic chemistry. *Angew. Chem. Int. Ed.* **2006**, *45*, 1844–1854. [[CrossRef](#)] [[PubMed](#)]
68. Bentley, T.W. Additivity rules using similarity models for chemical reactivity: Calculation and interpretation of electrofugality and nucleofugality. *Chem. A Eur. J.* **2006**, *12*, 6514–6520. [[CrossRef](#)] [[PubMed](#)]
69. Duynstee, E.F.J.; Grunwald, E. Organic Reactions Occurring in or on Micelles. II. Kinetic and Thermodynamic Analysis of the Alkaline Fading of Triphenylmethane Dyes in the Presence of Detergent Salts. *J. Am. Chem. Soc.* **1959**, *81*, 4542–4548. [[CrossRef](#)]
70. Duynstee, E.F.J.; Grunwald, E. Organic Reactions Occurring in or on Micelles. I. Reaction Rate Studies of the Alkaline Fading of Triphenylmethane Dyes and Sulfonphthalein Indicators in the Presence of Detergent Salts. *J. Am. Chem. Soc.* **1959**, *81*, 4540–4542. [[CrossRef](#)]
71. Valiente, M.; Rodenas, E. Benzene, toluene, and cyclohexane incorporation into hexadecyltrimethylammonium bromide micelles: Effect on the basic hydrolysis of crystal violet. *J. Colloid Interface Sci.* **1990**, *138*, 299–306. [[CrossRef](#)]
72. Leis, J.R.; Mejuto, J.C.; Peña, M.E. Comparison between the Kinetics of the Alkaline Fading of Carbocation Dyes in Water/Sodium Bis(2-ethylhexyl) Sulfosuccinate/Isooctane Microemulsions and in Homogeneous Media. *Langmuir* **1993**, *9*, 889–893. [[CrossRef](#)]
73. Cañadas, O.; Valiente, M.; Rodenas, E. Study of the cetyltrimethylammonium bromide/1,6-hexanediol/water system. *J. Colloid Interface Sci.* **1998**, *203*, 294–298. [[CrossRef](#)] [[PubMed](#)]
74. Mukherjee, L.; Mitra, N.; Bhattacharya, P.K.; Moulik, S.P. Kinetics in Microemulsion Medium. 4. Alkaline Fading of Crystal Violet in Aqueous (H<sub>2</sub>O/Aerosol OT/Isooctane and H<sub>2</sub>O/Aerosol OT/Decane) and Nonaqueous (Ethylene Glycol/Aerosol OT/Isooctane) Microemulsions. *Langmuir* **1995**, *11*, 2866–2871. [[CrossRef](#)]
75. García-Río, L.; Leis, J.R.; Mejuto, J.C.; Navarro-Vázquez, A.; Pérez-Juste, J.; Rodríguez-Dafonte, P. Basic hydrolysis of crystal violet in  $\beta$ -cyclodextrin/surfactant mixed systems. *Langmuir* **2004**, *20*, 606–613. [[CrossRef](#)] [[PubMed](#)]
76. Rakitin, A.R.; Pack, G.R. Monte Carlo calculations of ion distributions around micelles. *Colloids Surf. A Physicochem. Eng. Asp.* **2003**, *218*, 265–276. [[CrossRef](#)]
77. García-Río, L.; Leis, J.R.; Mejuto, J.C.; Pérez-Juste, J. Basic hydrolysis of m-nitrophenyl acetate in micellar media containing  $\beta$ -cyclodextrins. *J. Phys. Chem. B* **1998**, *102*, 4581–4587. [[CrossRef](#)]

78. Rodenas, E.; Perez-Benito, E.; Valiente, M. Effect of the location of reactants in the L<sub>2</sub> phase on the chemical reactivity. *Tenside Surfactants Deterg.* **1994**, *31*, 168–170. [[CrossRef](#)]
79. Schoental, R. Carcinogenic action of diazomethane and of nitroso-N-methyl urethane. *Nature* **1960**, *188*, 420–421. [[CrossRef](#)] [[PubMed](#)]
80. Astray, G.; Cid, A.; García-Río, L.; Hervella, P.; Mejuto, J.C.; Pérez-Lorenzo, M. Organic reactivity in AOT-stabilized microemulsions. *Prog. React. Kinet. Mech.* **2008**, *33*, 81. [[CrossRef](#)]
81. García-Río, L.; Mejuto, J.C.; Pérez-Lorenzo, M.; Rodríguez-Dafonte, P. Nitroso group transfer reactions between N-methyl-N-nitroso-P-toluene sulfonamide and N-alkylamines in CTAC1 micellar aggregates. *Prog. React. Kinet. Mech.* **2006**, *31*, 129–138. [[CrossRef](#)]
82. Fernández, I.; García-Río, L.; Hervés, P.; Mejuto, J.C. β-Cyclodextrin-micelle mixed systems as a reaction medium. Denitrosation of N-ethyl-N-nitroso-p-toluene-sulfonamide. *J. Phys. Org. Chem.* **2000**, *13*, 664–669. [[CrossRef](#)]
83. García-Río, L.; Hervés, P.; Mejuto, J.C.; Pérez-Juste, J.; Rodríguez-Dafonte, P. Pseudophase approach to the transfer of the nitroso group in water/AOT/SDS/isooctane quaternary microemulsions. *Langmuir* **2000**, *16*, 9716–9721. [[CrossRef](#)]
84. García-Río, L.; Hervés, P.; Leis, J.R.; Mejuto, J.C.; Pérez-Juste, J. Hydrolysis of N-methyl-N-nitroso-p-toluenesulphonamide in micellar media. *J. Phys. Org. Chem.* **1998**, *11*, 584–588. [[CrossRef](#)]
85. García-Río, L.; Leis, J.R.; Mejuto, J.C.; Pérez-Juste, J. Investigation of micellar media containing β-cyclodextrins by means of reaction kinetics: Basic hydrolysis of N-methyl-N-nitroso-p-toluenesulfonamide. *J. Phys. Chem. B* **1997**, *101*, 38. [[CrossRef](#)]
86. Bravo, C.; Hervés, P.; Leis, J.R.; Peña, M.E. Micellar effects in the acid denitrosation of N-Nitroso-N-methyl-p-toluenesulfonamide. *J. Phys. Chem.* **1990**, *94*, 8816–8820. [[CrossRef](#)]
87. García-Río, L.; Hervés, P.; Mejuto, J.C.; Parajó, M.; Pérez-Juste, J. Association Constant of Crystal Violet in Micellar Aggregates: Determination by Spectroscopic Techniques. *J. Chem. Res. Part S* **1998**, *11*, 716–717. [[CrossRef](#)]